

UNIVERSITY OF GOTHENBURG

This article has been accepted for publication in Monthly Notices of the Royal Astronomical Society ©: 2014 The Authors. Published by Oxford University Press on behalf of Royal Astronomical Society. All rights reserved.

Citation for the published paper:

Bakala, P.; Torok, G.; Karas, V.; Dovciak, M.; Wildner, M.; Wzientek, D.; Sramkova, E.; Abramowicz, Marek A; Goluchova, K.; Mazur, G. P.; Vincent, F. H. Power density spectra of modes of orbital motion in strongly curved space-time: obtaining the observable signal Monthly Notices of the Royal Astronomical Society, 439 (2) pp. 1933-1939 http://dx.doi.org/10.1093/mnras/stu076



MNRAS 439, 1933-1939 (2014) Advance Access publication 2014 February 12

Power density spectra of modes of orbital motion in strongly curved space-time: obtaining the observable signal

P. Bakala,^{1*} G. Török,¹ V. Karas,² M. Dovčiak,² M. Wildner,^{1,3} D. Wzientek,¹ E. Šrámková,¹ M. Abramowicz,^{1,4,5} K. Goluchová,¹ G. P. Mazur^{5,6} and F. H. Vincent^{5,7}

¹Institute of Physics, Faculty of Philosophy and Science, Silesian University in Opava, Bezručovo nám. 13, CZ-74601 Opava, Czech Republic ²Astronomical Institute, Boční II 1401, CZ-14131 Prague, Czech Republic

³Department of Theoretical Physics and Astrophysics, Faculty of Science, Masaryk University, CZ-61137 Brno, Czech Republic

⁴Physics Department, Gothenburg University, SE-412-96 Göteborg, Sweden

⁵Copernicus Astronomical Center, ul. Bartycka 18, PL-00-716 Warszawa, Poland

⁶Institute for Theoretical Physics, University of Warsaw, Hoza 69, PL-00-681 Warsaw, Poland

⁷Laboratoire AstroParticule et Cosmologie, CNRS, Universite Paris Diderot, 10 rue Alice Domon et Leonie Duquet, F-75205 Paris Cedex 13, France

Accepted 2014 January 10. Received 2013 November 21; in original form 2013 April 26

ABSTRACT

High-frequency quasi-periodic oscillations (HF QPOs) appear in the X-ray variability of several accreting low-mass binaries. In a series of works, it was suggested that these OPOs may have connection to inhomogeneities orbiting close to an inner edge of the accretion disc. In this paper, we explore the appearance of an observable signal generated by small radiating circular hotspots moving along quasi-elliptic trajectories close to the innermost stable circular orbit in the Schwarzschild space-time. Our consideration takes into account the capabilities of observatories that have been operating in the past two decades represented by the Rossi X-ray Timing Explorer (RXTE) and the proposed future instruments represented by the Large Observatory for X-ray Timing (LOFT). For these purposes, we choose such model parameters that lead to light curves comparable to those observed in Galactic black hole sources, in particular the microquasar GRS 1915+105. We find that when a weak signal corresponding to the hotspot Keplerian frequency is around the limits of the RXTE detectability, the LOFT observations can clearly reveal its first and second harmonics. Moreover, in some specific situations the radial epicyclic frequency of the spot can be detected as well. Finally, we also compare the signal produced by the spots to the signal produced by axisymmetric epicyclic disc-oscillation modes and discuss the key differences that could be identified via the proposed future technology. We conclude that the ability to recognize the harmonic content of the signal can help to distinguish between the different proposed physical models.

Key words: accretion, accretion discs – black hole physics – gravitation – methods: numerical-X-rays: binaries.

1 INTRODUCTION

Several low-mass X-ray binaries exhibit in the high-frequency part of their X-ray power density spectra (PDS) distinct peaks, so-called quasi-periodic oscillations (QPOs). The oscillations with the highest frequencies are sometimes displayed in the range of hundreds of Hz (HF QPOs). In neutron star (NS) sources, two simultaneous strong HF QPO peaks are often found (referred to as the upper and the lower QPO). The frequencies of these twin-peak QPOs are varying over time, sometimes by several hundred Hz on time-scales of days, and the fastest documented changes of twin QPO frequency

are of the order of Hz s⁻¹ (Paltani et al. 2004; Barret et al. 2005; Barret & Vaughan 2012). The frequency ratio sampled from the available observations of individual NS QPO sources often clusters (typically around the 3:2 value). The clustering can account either to incomplete data sampling, weakness of the two QPOs outside the limited frequency range, or to the intrinsic source clustering (see Abramowicz et al. 2003b; Belloni, Méndez & Homan 2005; Belloni et al. 2007; Barret & Boutelier 2008; Török et al. 2008a,b,c; Török 2009; Boutelier et al. 2010; Wang et al. 2013, for details). The HF QPOs observed in microquasars exhibit the 3:2 frequency ratio (e.g. Abramowicz & Kluźniak 2001; McClintock & Remillard 2006) as well. These QPOs are, however, somewhat different from the case of NS observations. Their frequencies are much more stable while their amplitudes are lower. One should also note that they are

^{*}E-mail: pavel.bakala@fpf.slu.cz

not displayed frequently and exhibit spectral differences from the NS QPOs (e.g. van der Klis 2006).

So far, there is no consensus on the QPO origin, and numerous models have been proposed (e.g. Alpar & Shaham 1985; Lamb et al. 1985; Miller, Lamb & Psaltis 1998; Psaltis et al. 1999; Wagoner 1999; Abramowicz & Kluźniak 2001; Wagoner, Silbergleit & Ortega-Rodríguez 2001; Titarchuk & Wood 2002; Rezzolla, Yoshida & Zanotti 2003; Pétri 2005; Zhang 2005; Kato 2007; Šrámková, Torkelsson & Abramowicz 2007; Stuchlík et al. 2008; Horák et al. 2009; Mukhopadhyay 2009; Lai et al. 2012, and others). Several but not all of the models expect that the generic QPO mechanism is the same for both NS and BH sources. Commonly, there is a belief that the HF OPOs carry important information about the inner accreting region dominated by the effects of strong Einstein's gravity. In several works, it was suggested that the observed modulation of X-ray flux may have some connection to the inhomogeneities propagating close to the inner edge of the accretion discs. This idea was first discussed within the hypothesis of PDS continuum origin due to the presence of spots on the accretion disc surface (e.g. Abramowicz et al. 1991; Mineshige, Takeuchi & Nishimori 1994).¹

Spots as the origin of QPOs have been discussed, in particular, in a series of papers (Stella & Vietri 1998a,b, 1999, 2002; Morsink & Stella 1999) that explain QPOs as a direct manifestation of modes of relativistic epicyclic motion of blobs at various radii *r* in the inner parts of the accretion disc. Within the model, the two observed HF QPO peaks arise due to the Keplerian and periastron precession of the relativistic orbits. In a similar manner, Čadež Calvani & Kostić (2008), Kostić et al. (2009), and Germana et al. (2009) introduced a concept in which the QPOs were generated by a tidal disruption of large accreting inhomogeneities. Recently, Germana (2012) argued that the behaviour of the azimuth phase $\phi(t)$ for non-closed quasielliptic orbits in the curved space–time can be responsible for the observed pairs of HF QPOs.

In this paper, we investigate the behaviour of the observable signal produced by radiating small circular hotspots. We discuss the detectability of the produced signal propagated from the strongly curved space-time region. In our discussion, we consider the capabilities of the present X-ray observatories represented by *Rossi X-ray Timing Explorer (RXTE)*, as well as the proposed future instruments represented by the *Large Observatory for X-ray Timing* (*LOFT*). We also compare the signal produced by spots to the signal obtained from another specific kind of simulations assuming axisymmetric epicyclic disc-oscillation modes.

2 TRACING THE SIGNAL FROM SPOT

Effects of special and general relativity act on photons emitted by the hot matter that orbits in the inner part of the accretion disc with velocity reaching values of percentages of speed of light. Complex fast calculations of the flux propagation through curved spacetime towards a distant observer represent a demanding, although, in principle, well-understood problem with practical astrophysical applications. In past decades, serious effort has been invested to solve it using mathematically elegant approaches, as well as straightforward ray-tracing methods profiting from currently available high computational power. Overall, various alternative strategies have been developed by different authors (e.g. de Felice, Nobili & Calvani 1974; Cunningham 1975; Laor 1991; Karas & Bao 1992; Karas, Vokrouhlicky & Polnarev 1992; Stuchlík & Bao 1992; Matt, Fabian & Ross 1993; Viergutz 1993; Zakharov 1994; Bromley, Chen & Miller 1997; Dabrowski et al. 1997; Bursa et al. 2004; Bakala et al. 2007; Vincent, Gourgoulhon & Novak 2012, and others). Particular application to the timing of radiating spots orbiting around black holes was discussed in greatest detail by Schnittman & Bertschinger (2004a,b) who elaborated the model of small radiating spots described as isotropic, monochromatic emitters following non-circular geodesic trajectories and also investigated effects given by large azimuthal shearing of such spots.

The expected spot-signal can be in principle used to model the response of X-ray detectors such as the proportional counter array on board the RXTE (Jahoda et al. 1996, 2006). The actual response of the instrument, however, depends on the complex time-dependent spectral behaviour of the whole source flux. In order to obtain relevant results one must then use a complete physical model of the accreting source radiation. First steps in this direction were taken in the above-mentioned work of Schnittman & Bertschinger who investigated their hotspot light curves assuming steady state disc models. This option for modelling the observable signal might be, however, difficult. The current observed radiation of Galactic microquasars exhibits a complex behaviour of transitions between particular X-ray states which are yet poorly understood, and the detailed self-consistent picture of full accreting system physics is rather missing so far. Another option which we follow here is to use an empirical model of the global source flux and add just the particular investigated model for the HF QPO variability including the predicted time-dependent energy spectra. This allows us to reproduce well the detectable signal using the response matrices of the RXTE and compare them to the results that can be achieved through the Large Area Detector currently proposed within the future LOFT mission (Feroci et al. 2012).

2.1 Simulation setup

Next, we investigate the signal from the motion of a small radiating spot obtained using the extended KYSPOT code (Dovčiak, Karas & Yaqoob 2004). For simplicity, we choose the zero black hole spin. We assume radial perturbations of the circular orbit of the radiating spot at the specific orbital radius $r_{3:1} = 6.75M$. At this radius, the radial epicyclic frequency v_r is exactly three times smaller than the Keplerian frequency v_k . The periastron precession frequency v_p then reaches the value $v_p = v_k - v_r = 2/3v_k$. The intrinsic local spot spectrum is chosen in a power-law form,

$$N(E) = kE^{-1.9}.$$
 (1)

For the further elaboration of observable effects, we set the black hole mass $M \doteq 11 \,\mathrm{M_{\odot}}$. The considered mass scales the Keplerian frequency of the spot to 160 Hz, close to one of HF QPO frequencies observed in the microquasar GRS 1915+105 (e.g. McClintock & Remillard 2006). For the spot trajectory we assume constant angular velocity equal to the Keplerian value $\Omega = \Omega_{\mathrm{K}}^{3:1}$ and two different amplitudes \mathcal{E} of the radial epicyclic oscillation - $\mathcal{E} = 0.1M$ and $\mathcal{E} = 1M$, respectively. These amplitudes correspond to a dimensionless pseudo-eccentricity

$$e \equiv \frac{(r_{\max} - r_{\min})}{(r_{\max} + r_{\min})} = \mathcal{E}/r$$
⁽²⁾

¹ In the same context, Abramowicz et al. (1992) argue that the surfaces of accretion discs must be populated by vortices. They point out that in differentially rotating fluids, vortices are not to be quickly destroyed by shear. Instead, they may be a very long lived objects, that survive in the disc like the Red Spot in Jupiter's atmosphere.

reaching the value of $e \doteq 0.01$ and $e \doteq 0.1$, respectively. The integration interval of the spot observation measured by an observer at the infinity is chosen as 10 s. Due to the technical limitations of the applied code, we restrict ourselves to this integration time in all further simulations presented in this paper.

We assume the global source flux described by the spectral distribution N(E) and power density spectra P(v),

$$N(E) = kE^{-2.5}, (3)$$

$$P(\nu) = p_0 \nu^{p_1} + \frac{1}{\pi} \frac{p_3 p_4}{(\nu - p_2)^2 + p_3^2},$$
(4)

where *k* is chosen to normalize the assumed count rate roughly to 1 Crab and $p_i = [0.001, -1.3, 2.5, 0.8, 0.002]$. This setup mimics the so-called high-steep power-law state in GRS 1915+105, including steep spectrum and power-law dominated variability with an additional broad Lorentzian component at low frequencies (McClintock & Remillard 2006).

2.2 Signal behaviour

Since the signal from the spot strongly depends on the source inclination i, we compare the results for two representative values of i corresponding to the nearly equatorial view and the view close to vertical axes.

2.2.1 Nearly equatorial view

In Fig. 1, we include amplitude spectra and time-dependent energy spectra of the net spot-signal calculated for the distant observer assuming $i = 80^{\circ}$. The spot-signal is dominated by the Keplerian frequency and its harmonics amplified by relativistic effects which is well illustrated by the amplitude spectrum on the left-hand panel of Fig. 1. The eccentricity given by e = 0.01 causes only a negligible modulation at the radial and precession frequency. The increased eccentricity corresponding to e = 0.1 can be well recognized in the amplitude spectra, but the signal is still dominated by the Keplerian frequency and its harmonics. The time-dependent energy spectra of the spot are depicted in the right-hand panel of Fig. 1. We can see that they clearly reveal the signatures of relativistic redshift effects.

So far we have reproduced just the variability and spectra of the net spot flux. In order to assess the observable effects, we have to study the total composition of the net spot flux together with the global source flux given by equations (3) and (4). Assuming this composed radiation, we can consider the capabilities of the *RXTE* and *LOFT* instruments using their response matrices and provided software tools. The time-dependent spectra describing the composed radiation are then convolved with the appropriate response matrix giving an estimate of the observed data. These are Fourier transformed to the resulting power spectra. The detectability of the spot signatures then depends obviously on the fraction of photons from the spot in the total flux. Hereinafter, we refer to this quantity which determines the signal-to-noise ratio shortly as signal fraction n.

Fig. 2 shows PDS resulting from the *RXTE* and *LOFT* simulations assuming various levels of *n* and the inclination $i = 80^{\circ}$ (i.e. nearly equatorial view). It includes the cases when the signal is weak for the *RXTE* and there are no significant features within its PDS as well as the high signal fraction when first two harmonics of the Keplerian frequency can be seen. Comparing both panels of this figure, we can deduce that when the weak QPO signal corresponding to the hotspot Keplerian frequency is around the limits of the *RXTE* detectability, the *LOFT* observations will clearly reveal its first and second harmonics. We checked that there is, in practice, no qualitative difference between the cases of e = 0.01 and e = 0.1. It is therefore unlikely that the periastron precession or radial epicyclic frequency can be detected in addition to the harmonics when the inclination angle is close to the equatorial plane.

2.2.2 View close to vertical axes

For $i = 30^{\circ}$, the signal is dominated by the Keplerian frequency but the harmonics are much less amplified in comparison to the nearly equatorial view (see the left-hand panel of Fig. 3). The e = 0.01pseudo-eccentricity again causes rather negligible modulation at the radial and the precession frequency. Nevertheless, we can see that the increased eccentricity of e = 0.1 affects the variability more than for the large inclination angle. Furthermore, the modulation at the radial frequency is comparable to those at the second harmonics of the Keplerian frequency. The time-dependent energy spectra are depicted in the right-hand panel of Fig. 3.

Fig. 4 includes PDS resulting from *RXTE* and *LOFT* simulations assuming various levels of n. It is drawn for e = 0.1 and includes the few cases when the signal is weak for the *RXTE* and there are no significant features within its PDS, plus one case when some feature at the Keplerian frequency can be seen. Comparing both panels of this figure, one can deduce that when the weak QPO signal



Figure 1. Expected net spot flux measured by a distant observer close to the equatorial plane. Left: amplitude spectrum. Right: time-dependent energy spectra drawn for a distant observer.



Figure 2. Leahy-normalized PDS obtained for various levels of signal fraction *n* assuming nearly equatorial view ($i = 80^{\circ}$) and pseudo-eccentricity e = 0.1. The colour-coded flags emphasize the signal excess above the noise level for given *n*. Left: outputs for an exemplary source regarding the *RXTE* capabilities. Right: outputs for the same source luminosity considering the *LOFT* capabilities. Note that the lowest displayed values of *n* illustrated by the blue and yellow colour do not indicate any significant features within the *RXTE* PDS. On the other hand, the *LOFT* PDS reveals the Keplerian frequency (blue and yellow PDS), respectively, its first two harmonics (yellow PDS).



Figure 3. Expected net spot flux measured by a distant observer close to vertical axes. Left: amplitude spectrum. Right: time-dependent energy spectra drawn for the distant observer.



Figure 4. Leahy-normalized PDS obtained for various levels of signal fraction *n* assuming a view close to the vertical axes ($i = 30^{\circ}$) and the pseudo-eccentricity e = 0.1. The colour-coded flags emphasize the signal excess above the noise level for given *n*. Left: outputs for an exemplary source, considering the *RXTE* capabilities. Right: outputs for the same source luminosity, considering the *LOFT* capabilities. Note that *RXTE* PDS includes a barely significant excess of power at 160 Hz only for the highest displayed value of signal fraction *n*. For the same value, the *LOFT* PDS reveals the first two harmonics and also the radial epicyclic frequency, $v_r = 53$ Hz. Moreover, combinational harmonic frequency, $4v_r = 2v_P \doteq 213$ Hz, can be seen as well.



Figure 5. Left: amplitude distribution and its inclination dependence expected for the drifting spot observed from infinity. Right: illustration of the corresponding PDS drawn for the *RXTE* and *LOFT* observations and inclination $i = 65^{\circ}$.

corresponding to the hotspot Keplerian frequency is around the limits of the *RXTE* detectability, the *LOFT* observations can clearly reveal its first and second harmonics, and in addition the radial epicyclic frequency v_r and its combinational harmonics $4v_r = 2v_p$. Although it is not directly illustrated in the figure, it can be shown that decreasing eccentricity to e = 0.01 leads to a similar PDS, only without the features at v_r and $4v_r$.

3 BROAD PEAKS AND THEIR HARMONICS

In the previous simplified analysis, we assumed the single spot orbiting with the fixed Keplerian period. So far, there is no agreement on the physical origin and properties of the hypothetic clumps orbiting in the disc. In various proposed scenarios, the spot-signal (de)coherence is commonly related either to the radial distribution of several spots or to the continuous change of the spot orbital radius (see e.g. Karas 1999; Pecháček, Karas & Czerny 2008; Bini et al. 2012, for details and discussion). In the left-hand panel of Fig. 5, we show PDS resulting from consideration of spots created at r = 6.75M and drifted to r = 6M (ISCO). The figure illustrates well the strong inclination dependence of the ratio between the amplitudes observed at the Keplerian and harmonic frequencies. The example of instrument response to the total model flux is illustrated on the right-hand panel of Fig. 5 (the mass M is set to the same value as in the previous sections).

3.1 The 3:2 peaks produced by spots and tori

The radial distribution or drifting of the spots can clearly result in various levels of signal coherence. Nevertheless, small circular spots related to a single preferred radius do not reproduce the often observed 3:2 frequency ratio. Motivated by Karas (1999), we also consider a somewhat more complicated arbitrary scheme where the multiple spots are created and drifted around radii close to two preferred orbits with Keplerian frequencies roughly in the 3:2 ratio. These orbits are set as $r_2 = 8M$ (radius where the radial epicyclic frequency reaches the maximum value) and $r_3 = 6M$ (ISCO). Spots are then created within the regions $[r_i + \delta r, r_i]$ with the size given by $\delta r = 0.75M$. The illustration of this scenario is shown in the left-hand panel of Fig. 6. Since we keep the same black hole mass as above, $M \doteq 11 \text{ M}_{\odot}$, our setup leads to the main observable frequencies around 110 and 160 Hz (see panels a and c of Fig. 7 drawn for signal fraction $n \doteq 10$ per cent and $i = 65^\circ$). In the following consideration, we compare the PDS obtained for this setup to the PDS resulting from the model of the oscillating optically thin torus slowly drifting through the resonant radius $r_{3:2}$.

For the torus kinematics, we assume the m = 0 radial and vertical oscillations with equal intrinsic amplitudes. The possible QPO origin in the resonances between this or similar disc-oscillation modes has been extensively discussed in works of Abramowicz & Kluźniak (2001), Abramowicz et al. (2003a,b), Kluźniak et al. (2004), Horák (2005), Török et al. (2005), and several other authors. Here, we adopt the concept previously investigated by Bursa et al. (2004) who focused on optically thin torus with slender geometry.

The visual appearance of torus influenced by lensing and Doppler effects is illustrated in the right-hand panel of Fig. 6. The periodic changes of the observed luminosity are partially governed by the radial oscillations due to changes of the torus volume while the vertical oscillations modulate the flux just due to lensing effects in the strong gravitational field. The contribution of the two individual oscillations to the variations of the observed flux thus strongly depends on the inclination angle (see also Mazur et al. 2013). Here, we set $i = 65^{\circ}$ where the fractions of the power in the two observed peaks are comparable. We set the black hole mass $M = 5.65 \text{ M}_{\odot}$ and a = 0 ($r_{3:2} = 10.8M$), implying that the two oscillatory frequencies are $v_{\theta}(r_{3:2}) = 160 \text{ Hz}$ and $v_{r}(r_{3:2}) \doteq 110 \text{ Hz}$. Assuming this setup, we produce torus drift light curves for the interval $r/r_{3,2} \in [0.97,$ 1.03]. The resulting PDS drawn for signal fraction $n \doteq 10$ per cent is included in Fig. 7 (panels b and d). We note that similar PDS can be reached assuming e.g. a near extreme rotating black hole with a = 0.98 and $M \doteq 18$ M_{\odot}.

Using Fig. 7, we can finally confront the predictions for spots drifted around preferred radii to those expected for the oscillating torus slowly passing the resonant orbit $r_{3:2}$. Inspecting this figure, we can find that the *RXTE* PDS obtained for the given setup of the two models are rather similar. On the other hand, the *LOFT* PDS reveal the presence/absence of the harmonics in addition to the 3:2 peaks.

3.2 Other examples of the harmonic signature of spot motion

Higher harmonics should appear in the signal generated by small spots in various situations. In the following, we justify their possible detectability assuming two more examples of the spot motion.

Panels (e) and (g) of Fig. 7 marked as Spots-S include PDS resulting for the group of 200 stationary spots ($i = 65^{\circ}$). These spots were spatially distributed around the two orbits r_2 and r_3



Figure 6. Left: sketch of the 3:2 QPO model scheme based on spots orbiting close to two preferred radii. Right: illustration of the role of lensing and Doppler effects in the visual appearance of torus located at the resonant orbit $r_{3:2}$.



Figure 7. Panels (a)–(d): comparison between the multiple spot and oscillating torus PDS obtained for the two instruments. Superimposed red curves indicate various multi-Lorentzian models. Panels (e)–(h): results of simulations assuming stationary spots distributed in space (Spots-S) and spots with short lifetimes located just at the two orbits (Spots-T).

assuming a superposition of Gaussian radial distributions of deviations $\sigma_{2,3} = (0.3, 0.1)$ and random orbital phases. In contrary, panels (f) and (h) of the same figure marked as Spots-T assume a non-stationary situation with spots that repetitively appear only at the two preferred radii r_2 and r_3 ($i = 65^\circ$). The spots were created with long delays with respect to their random lifetimes. These were much shorter than the integration time of the simulation. Apparently, in both chosen examples, we can find harmonics representing the signature of the spot motion.

Inspecting the Spot-S and Spot-T situations, we have found clear harmonics in the expected amplitude spectra for any choice of the model parameters. Nevertheless, we note that their detectability in the observational PDS was not high in all cases. Moreover, as illustrated in the case of Spots-T, various models can produce strong noise at low frequencies.

Detailed elaboration of the question which setup guarantees the detectability of the harmonics thus requires a full study of the related (rich) parameter space. We do not intend to make such a study in this work, but it should be discussed elsewhere.

4 DISCUSSION AND CONCLUSIONS

We identified the signatures of the spot motion mostly with the harmonic content of the observable signal. Remarkably, for large inclination angles, the *LOFT* observations could easily reveal the Keplerian frequency together with its first and second harmonics when the strongest (but still very weak) single signal is around the limits of the *RXTE* detectability. Nevertheless, the radial epicyclic frequency could be also found providing that the inclination is small. In our analysis, we have paid attention to the timing signatures of the motion of small circular spots radiating isotropically from the slightly eccentric geodesic orbits. The case of highly eccentric orbits and/or spot having large azimuthal shear will be presented elsewhere.

The comparison between spot and torus models in Section 3 has been obtained for specific kinematic models. Its general validity is thus limited. For instance, a consideration of resonance-driven effects or torus geometrical thickness could also give rise to some harmonic content in the signal from the oscillating tori. Despite

ACKNOWLEDGEMENTS

We acknowledge the support of the Czech grant GAČR 209/12/P740, Polish grant NCN UMO-2011/01/B/ST9/05439, and the Swedish VR grant. We further acknowledge the project CZ.1.07/2.3.00/20.0071 'Synergy' aimed to support international collaboration of the Institute of physics in Opava, and the internal SU student grant SGS/11/2013. Part of the work was supported by the COST-CZ project MP1304 (Astrophysics of toroidal fluid structures near compact stars) in the Czech Republic, and the European Union Seventh Framework Programme under the grant agreement No. 312789. We thank to the anonymous referee for his comments and suggestions that helped to improve the paper.

REFERENCES

- Abramowicz M. A., Kluźniak W., 2001, A&A, 374, L19
- Abramowicz M. A., Bao G., Lanza A., Zhang X.-H., 1991, A&A, 245, 454
- Abramowicz M. A., Lanza A., Spiegel E. A., Szuszkiewicz E., 1992, Nature, 356, 41
- Abramowicz M. A., Karas V., Kluźniak W., Lee W. H., Rebusco P., 2003a, PASJ, 55, 466
- Abramowicz M. A., Bulik T., Bursa M., Kluźniak W., 2003b, A&A, 404, L21
- Alpar M. A., Shaham J., 1985, Nature, 316, 239
- Bakala P., Čermák P., Hledík S., Stuchlík Z., Truparová K., 2007, Cent. Eur. J. Phys., 5, 599
- Barret D., Boutelier M., 2008, New Astron. Rev., 51, 835
- Barret D., Vaughan S., 2012, ApJ, 746, 131
- Barret D., Kluźniak W., Olive J. F., Paltani S., Skinner G. K., 2005, MNRAS, 357, 1288
- Belloni T., Méndez M., Homan J., 2005, A&A, 437, 209
- Belloni T., Homan J., Motta S., Ratti E., Méndez M., 2007, MNRAS, 379, 247
- Bini D., Falanga M., Geralico A., Stella L., 2012, Classical Quantum Gravity, 29, 065014
- Boutelier M., Barret D., Lin Y., Török G., 2010, MNRAS, 401, 1290
- Bromley B. C., Chen K., Miller W. A., 1997, ApJ, 475, 57
- Bursa M., Abramowicz M. A., Karas V., Kluzniak W., 2004, ApJ, 617, L45
- Čadež A., Calvani M., Kostić U., 2008, A&A, 487, 527
- Cunningham C. T., 1975, ApJ, 202, 788
- Dabrowski Y., Fabian A. C., Iwasawa K., Lasenby A. N., Reynolds C. S., 1997, MNRAS, 288, L11
- de Felice F., Nobili L., Calvani M., 1974, A&A, 30, 111
- Dovčiak M., Karas V., Yaqoob T., 2004, ApJS, 153, 205
- Feroci M. et al., 2012, Exp. Astron., 34, 415
- Germana C., 2013, MNRAS, 430, L1
- Germana C., Kostič U., Čadež A., Calvani M., 2009, in Rodriguez J., Ferrando P., eds, AIP Conf. Proc. Vol. 1126, SIMBOL-X: Focusing on the Hard X-ray Universe. Am. Inst. Phys., New York, p. 367
- Horák J., 2005, Astron. Nachr., 326, 824
- Horák J., Abramowicz M. A., Kluźniak W., Rebusco P., Török G., 2009, A&A, 499, 535
- Jahoda K., Swank J. H., Giles A. B., Stark M. J., Strohmayer T., Zhang W., Morgan E. H., 1996, in Siegmund O. H., Gummin M. A., eds, Proc. SPIE, Vol. 2808, EUV, X-Ray, and Gamma-Ray Instrumentation for Astronomy VII. SPIE, Bellingham, p. 59
- Jahoda K., Markwardt C. B., Radeva Y., Rots A. H., Stark M. J., Swank J. H., Strohmayer T. E., Zhang W., 2006, ApJS, 163, 401
- Karas V., 1999, PASJ, 51, 317
- Karas V., Bao G., 1992, A&A, 257, 531
- Karas V., Vokrouhlicky D., Polnarev A. G., 1992, MNRAS, 259, 569
- Kato S., 2007, PASJ, 59, 451

- Kluźniak W., Abramowicz M. A., Kato S., Lee W. H., Stergioulas N., 2004, ApJ, 603, L89
- Kostić U., Čadež A., Calvani M., Gomboc A., 2009, A&A, 496, 307
- Lai D., Fu W., Tsang D., Horak J., Yu C., 2013, in Zhang C. M., Belloni T., Mendez M., Zhang S. N., eds, Proc. IAU Symp. 290, Feeding Compact Objects: Accretion on All Scales. Cambridge Univ. Press, Cambridge, p. 57
- Lamb F. K., Shibazaki N., Alpar M. A., Shaham J., 1985, Nature, 317, 681 Laor A., 1991, ApJ, 376, 90
- McClintock J. E., Remillard R. A., 2006, in Walter L., van der Klis M., eds, Cambridge Astrophysics Ser. 39, Compact Stellar X-ray Sources. Cambridge Univ. Press, Cambridge, p. 157
- Matt G., Fabian A. C., Ross R. R., 1993, MNRAS, 264, 839
- Mazur G. P., Vincent F. H., Johansson M., Šrámková E., Török G., Bakala P., Abramowicz M. A., 2013, A&A, 554, A57
- Miller M. C., Lamb F. K., Psaltis D., 1998, ApJ, 508, 791
- Mineshige S., Takeuchi M., Nishimori H., 1994, ApJ, 435, L125
- Morsink S. M., Stella L., 1999, ApJ, 513, 827
- Mukhopadhyay B., 2009, ApJ, 694, 387
- Paltani S., Barret D., Olive J. F., Skinner G. K., 2004, in Combes F., Barret D., Contini T., Meynadier F., Pagani L., eds, SF2A-2004: Semaine de l'Astrophysique Francaise. EdP-Sciences, Les Ulis Cedex, France, p. 381
- Pecháček T., Karas V., Czerny B., 2008, A&A, 487, 815
- Pétri J., 2005, A&A, 439, L27
- Psaltis D. et al., 1999, ApJ, 520, 763
- Rezzolla L., Yoshida S., Zanotti O., 2003, MNRAS, 344, 978
- Schnittman J. D., Bertschinger E., 2004a, ApJ, 606, 1098
- Schnittman J. D., Bertschinger E., 2004b, in Kaaret P., Swank J. H., Lamb F. K., eds, AIP Conf. Proc. Vol. 714, X-RAY TIMING 2003: Rossie and Beyond. Am. Inst. Phys., New York, p. 40
- Šrámková E., Torkelsson U., Abramowicz M. A., 2007, A&A, 467, 641
- Stella L., Vietri M., 1998a, in Paul J., Montmerle T., Aubourg E., eds, Abstr. 19th Texas Symp. on Relativistic Astrophysics and Cosmology. CEA, Saclay, France
- Stella L., Vietri M., 1998b, ApJ, 492, L59
- Stella L., Vietri M., 1999, Phys. Rev. Lett., 82, 17
- Stella L., Vietri M., 2002, in Gurzadyan V. G., Jantzen R. T., Ruffini R., eds, Proc. 9th Marcel Grossmann Meeting, On Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories. World Scientific, Singapore, p. 426 Stuchlík Z., Bao G., 1992, Gen. Relativ. Gravit., 24, 945
- Stuchlík Z., Konar S., Miller J. C., Hledík S., 2008, A&A, 489, 963
- Titarchuk L., Wood K., 2002, ApJ, 577, L23
- Török G., 2009, A&A, 497, 661
- Török G., Abramowicz M. A., Kluźniak W., Stuchlík Z., 2005, A&A, 436,
- Török G., Bakala P., Stuchlík Z., Čech P., 2008a, Acta Astron., 58, 1
- Török G., Abramowicz M. A., Bakala P., Bursa M., Horák J., Kluźniak W., Rebusco P., Stuchlík Z., 2008b, Acta Astron., 58, 15
- Török G., Abramowicz M. A., Bakala P., Bursa M., Horák J., Rebusco P., Stuchlík Z., 2008c, Acta Astron., 58, 113
- van der Klis M., 2006, in Lewin W. H. G., van der Klis M., eds, Compact Stellar X-Ray Sources. Cambridge Univ. Press, Cambridge, p. 39
- Viergutz S. U., 1993, A&A, 272, 355
- Vincent F. H., Gourgoulhon E., Novak J., 2012, Classical Quantum Gravity, 29, 245005
- Wagoner R. V., 1999, Phys. Rep., 311, 259
- Wagoner R. V., Silbergleit A. S., Ortega-Rodríguez M., 2001, ApJ, 559, L25
- Wang D. H., Chen L., Zhang C. M., Lei Y. J., Qu J. L., 2013, preprint (arXiv:1301.5478)
- Zakharov A. F., 1994, MNRAS, 269, 283
- Zhang C. M., 2005, Chin. J. Astron. Astrophys., 5, 21

This paper has been typeset from a TEX/LATEX file prepared by the author.