

Article Type: Original Article

Issue title:

‘Transgenerational Plasticity, Epigenetics and the Evolution of Marine Species Under Global Change’

Paper title:

‘Will Life find a Way? Evolution of Marine Species Under Global Change’

Piero Calosi^{1*}, Pierre De Wit², Peter Thor³, Sam Dupont⁴

¹ Département de Biologie, Chimie et Géographie, Université du Québec à Rimouski, Rimouski, QC, Canada

² Department of Marine Sciences, University of Gothenburg, Strömstad, Sweden

³ Framcentre, Norwegian Polar Institute, Tromsø, Norway

⁴ Department of Biological and Environmental Sciences, University of Gothenburg, Fiskebäckskil, Sweden

* corresponding author piero_calosi@uqar.ca

ABSTRACT

Projections of marine biodiversity and implementation of effective actions for its maintenance in face of current rapid global environmental change are constrained by our limited understanding of species' adaptive responses, including, transgenerational plasticity, epigenetics, natural selection. This special issue presents 13 novel studies, which employ experimental and modeling approaches to: (1) Investigate plastic and evolutionary responses of marine species to major global change drivers; (2) ask relevant broad eco-evolutionary questions, implementing well-designed multiple species and populations studies; (3) show the advantages of using advanced experimental designs and tools; (4) construct novel model organisms for marine evolution; (5) help identifying future challenges

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/eva.12418

This article is protected by copyright. All rights reserved.

for the field, and (6) highlight the importance of incorporating existing evolutionary theory into management solutions for the marine realm. What emerges is that at least some populations of marine species are able to adapt to future global change conditions. However, marine organisms' capacity for adaptation appears finite, due to evolutionary trade-offs and possible rapid losses in genetic diversity. This further corroborate the idea that acquiring an evolutionary perspective on how marine life will respond to the selective pressure of future global changes will guide us in better identifying which conservation efforts will be most needed, and most effective.

<<It is difficult to believe in the dreadful but quiet war lurking just below the serene facade of nature>> (Charles Darwin, 1859)

The chemical and physical evidence for on-going anthropogenic global change is now so prevalent that the conclusion that our climate is drastically changing is considered indisputable (IPCC 2013). On the other hand, and despite the tremendous effort by the Intergovernmental Panel on Climate Change (IPCC 2013) to synthesize our present understanding of the biological implications of global change, biological evidence corroborating the existence of ubiquitous mechanisms governing species' responses to future environmental challenges are somewhat lagging behind (Dupont and Pörtner 2013, Melzner et al. 2009, Kroeker et al 2013, Storch et al. 2014, Wittman and Pörtner 2015). This discrepancy has contributed preventing us from producing more conclusive projections on the fate of living systems under global change. What appears to be certain, it is that we are on the brink of a global biodiversity crisis (Barnowsky et al. 2011). It is thus unlikely that any extant species and ecosystem will be able to survive the on-going planetary environmental changes without actually changing. In fact, whilst migration can temporarily help prevent a species' global extinction, ultimately it is only through evolutionary adaptation that populations and species can be rescued from local and global extinction (Gonzalez et al. 2012). Nonetheless, phenotypic plasticity may buy additional time for adaptation to occur (Munday et al. 2013, Godbold and Calosi 2013, Reusch 2013, Sunday et al. 2014), and also provide a mechanism for adaptation to occur rapidly (Pigliucci et al. 2006, Ghalambor et al. 2015). Finally, extant levels of adaptation to local conditions may mediate populations' sensitivity to future global change drivers (e.g. Lardies et al. 2014, Wood et al. 2016). For these reasons, the investigation of populations' and species' ability to mount plastic and adaptive responses to prevalent environmental changes is an absolute priority (e.g. Pespeni et al. 2013), if we are to identify which populations, species and assemblages will survive global change, and which are more likely to go extinct (Hall-Spencer et al. 2008, Calosi et al. 2013, Lucey et al. 2015). As current efforts have to a large extent focused on individual species' abilities to cope with short-term changes through plastic responses, in order to

make critical predictions of long-term responses, it is essential to gain an understanding of the mechanisms behind the complex interactions between plasticity, evolution and non-genetic inheritance (epigenetics).

The study of evolution is one of the central themes of modern biology (Darwin 1859, Dobzhansky 1951, Huxley 1952, Dobzhansky 1973, Margulis 1998, Nobel 2015). Species' capacity to mount evolutionary responses to fluctuations and changes in the environment has been investigated for decades both *via* comparative (see in Hochachka and Somero 2002, Stillman and Paganini 2015) and correlative methods (see Colin and Dam 2002, Gaston et al. 2009, Bozinovic et al 2011, Dam 2013). More recently, in order to overcome some of the limitations of these former methods the implementation of experimental evolutionary methods has been favoured (e.g. Bennett et al. 1992, Kellerman et al. 2009, Garland and Rose 2009). Nonetheless, in the field of Marine Global Change Biology the investigation of the capacity of biological systems to adapt to the on-going rapid environmental change has been largely overlooked, at least until very recently (Munday et al. 2013, Godbold and Calosi 2013, Reusch 2013, Sunday et al. 2014). This situation may result from the historical lack of true marine model systems, particularly for multicellular organisms when compared to terrestrial systems (e.g. *Drosophila*, *Arabidopsis*). In part, this has been a consequence of the difficulty of working with long-lived species, in a poorly understood environment, as well as having to deal with maintaining desired environmental conditions in sea water in a laboratory. Nonetheless, as marine systems, just like other biological systems are intrinsically plastic (Ghalambor et al. 2007) and have the ability to evolve (Darwin 1859), in some cases rapidly (e.g. Thor and Dupont 2015, Ghalambor et al. 2015), these features can no longer be ignored when trying to project the responses of marine populations, species and assemblages to rapid changes in multiple environmental drivers.

There is no doubt that the IPCC (2013) has generated an in-depth synthesis of the patterns through which marine species and ecosystems presently respond to the on-going global change and may do so in the future (Pörtner et al. 2014). If we are to critically improve current predictions of the fate of global biodiversity under the current environmental change, advances in understanding of the drivers and mechanisms behind marine evolution are required. In this sense, the investigation of trans-generational plastic and evolutionary responses of fitness-related traits under global change scenarios, and the identification of the underpinning physiological genetic and non-genetic mechanisms, is central to advance our current understanding of how marine organisms will be able to cope with future environmental challenges. Using an evolutionary approach will help us avoid potential overestimations or underestimations of the biological implications of global change (Dam 2013).

Consequently, this special issue aims to collect novel, cutting-edge studies, which represent a further proof for the idea that the investigation of evolution within the context of marine global change is imperative, and can help guiding environmental management and conservation solutions under the ongoing rapid global change.

The specific objectives of this special issue are to: (1) Investigate plastic and evolutionary responses of ecologically important marine species to some of the major global change drivers (e.g. ocean warming, ocean acidification, salinity changes), both as single drivers but also combined (simultaneous and sequential); (2) move toward asking relevant broad eco-evolutionary questions, implementing well-designed multiple species and populations studies; (3) show the advantages of using advanced experimental designs and appropriate tools (from high-throughput DNA sequencing and novel methods for studying methylation patterns, to mathematical modelling); (4) move beyond current limitations by constructing novel model organisms for evolution in the marine realm; (5) help identify some of the future challenges for the field of marine global change biology, and finally (6) highlight the importance of incorporating existing evolutionary theory into management solutions for the marine realm.

This special issue consists of thirteen original manuscripts, focusing on unicellular organisms, macroalgae, invertebrates, and vertebrates as study models. Most importantly, these works cover a broad range of approaches and topics within the context of future global change scenarios. These include (i) the investigation of the significance of local adaptation in defining populations' responses; (ii) the importance of trans- and multigenerational responses to mediate species' plastic responses; (iii) the possibility for rapid evolution to occur; and (iv) the relevance of epigenetic mechanisms, as well as evolutionary trade-offs, in mediating species' responses. From these studies a number of relevant messages and lessons have emerged, and are briefly summarised below.

Local Adaptation

A species' level of local adaptation, here defined as the process of evolution of a given population in response to the prevalent local environmental regimes (Williams 1966) in the face of gene flow from nearby populations, will be critical in define populations' responses to future environmental conditions, by either providing a buffer for future negative impacts, or increasing sensitivity levels (e.g. Sanford et al. 2011, Calosi et al. 2013, Dam 2013, Pespeni et al. 2013, Savolainen 2013). Within this special issue, Padilla-Gamino et al. (2016) have used multiple life stages of different species of coralline algae to test the hypothesis that populations living in habitats characterised by higher variability and elevated levels of seawater $p\text{CO}_2$ will be less affected by future ocean acidification, when compared to

populations from habitats characterised by more stable and low levels of seawater $p\text{CO}_2$. They were able to show that spores are less sensitive to elevated $p\text{CO}_2$ than adults, and found more marked impacts in populations found in habitats characterised by lower variability and lower levels of seawater $p\text{CO}_2$. These findings have important implications for the conservation of these important ecosystem engineers in the future ocean. Indeed, the investigation of local adaptation must become a conservation and resource management priority (Lucey et al. 2016).

On the other hand, Lucey et al. (2016) carried out a reciprocal transplant on individuals of the sessile calcifying polychaete *Simplaria* sp. from a population inhabiting a naturally elevated $p\text{CO}_2$ volcanic vent area, and a population from a nearby control area exposed to unaltered water chemistry conditions. Their results indicate that in this taxon neither local adaptation nor phenotypic plasticity may suffice to buffer the negative impacts of future ocean acidification. In more detail, Lucey et al. (2016) showed that regardless of their original environmental conditions, both populations showed low fitness levels, increased tube growth rates and similar plastic responses when exposed to elevated $p\text{CO}_2$ conditions, suggesting that local adaptation to a low pH environment had not occurred, and thus that long-term exposure had not caused any substantial phenotypic changes.

Results from these two studies suggest that local adaptation to future conditions may not be a ubiquitous process in the marine environment. Large variability in evolutionary and plastic responses may exist, most likely resulting from differences in life-history strategies, population size, fecundity and gene flow. Understanding the relative contributions of these parameters to local adaptive capability will enable us to widen our knowledge on the importance of the process of adaptation to counter environmental change, and ultimately use it to promote the conservation of marine biodiversity. Finally, Padilla-Gamino et al. (2016) and Lucey et al. (2016) both show the value of comparing populations living under differing environmental regimes as an approach to study marine organisms potential for evolution under global change.

Trans-generational and Multigenerational Studies, and Evidence for Rapid Selection

Trans-generational effects, defined as changes in offspring phenotype due to stress-exposure of the parental generation have the potential to buffer species against environmental changes (Sunday et al. 2014). In this special issue, Donelson et al. (2016) use a model coral reef fish (*Acanthochromis polyacanthus*) to investigate the impact of different heat exposure of parents on the next generation's reproductive output ability and the quality of offspring produced. Interestingly, they found that a gradual warming over two generations resulted in greater plasticity of the reproductive traits investigated, when compared to fish that experienced the same increase within one generation. Similarly, evidence for positive effect of trans-generational exposure in helping re-stabilising

reproductive output levels following a rapid change in $p\text{CO}_2$ are also provided by Rodriguez-Romero et al. (2015), using a laboratory strain of an emerging marine polychaete model (*Ophryotrocha labronica*). These studies (Donelson et al. 2016, Lucey et al. 2016) suggest that trans-generational plasticity can induce full restoration of fitness-related traits, which may not be observed with developmental plasticity alone. Furthermore, Rodriguez-Romero et al. (2015) also conducted a mutual transplant experiment, following seven generations of exposure to differing $p\text{CO}_2$ conditions, providing evidence for the possible occurrence of rapid adaptation in a marine organism to a rapid environmental change. Rodriguez-Romero et al. (2015) show the importance of conducting multi-generational experiments in order to provide more realistic estimates for marine metazoans' responses to future environmental changes. However, they also highlight the limitations of interpreting the evolutionary significance of the outcome of transgenerational and multigenerational experiments, without the use of physiological and genetic tools, often not available for non-model organisms.

A number of studies in this special issue integrate novel physiological and genetic tools in the investigation of marine organisms' responses to global change drivers. For example, Shama et al. (2016) investigated differences in mitochondrial respiratory capacity and gene expression across three generations in marine sticklebacks exposed to heat stress, either in an acute fashion or through development, allowing for some acclimation to occur. They used an advanced cross-breeding experimental design, and demonstrated that the mechanisms underlying trans-generational effects persist across multiple generations, leading to phenotypes for mitochondrial respiratory capacity and gene expression that depend on both the type of acclimation and the environmental mismatch between generations. In addition, De Wit et al. (2015) further corroborated the evolutionary significance of the mitochondrial function in underpinning species' transgenerational responses to global changes. In order to do this, they exposed specimens of the copepod *Pseudocalanus acuspes* to different $p\text{CO}_2$ conditions over two successive generations, followed by a reciprocal transplant experiment (Thor and Dupont 2015). After this, they used a physiological hypothesis-testing strategy to mine both gene expression and nucleotide sequence data showing that exposure to elevated $p\text{CO}_2$ appears to impose selection in copepods on both mitochondrial and ribosomal function, and that these changes might be related to changes in RNA transcription activity. The important consequence of this work is that De Wit et al. (2015) show that evolution of fitness-related traits can occur rapidly in marine metazoans exposed to future global change scenarios, especially in species with high standing genetic variation and large population sizes. This gives some hope that selection acting on existing phenotypic and genetic diversity can promote the rescue of some marine metazoans within the context of future global change conditions (Munday et al. 2013, Reusch 2013, Sunday et al. 2014).

Genetic diversity could rapidly diminish in the face of rapid environmental changes, as shown by Lloyd et al. (2016) in the larvae of the purple sea urchin (*Strongylocentrotus purpuratus*) exposed to elevated $p\text{CO}_2$ conditions. Lloyd et al. (2016) showed a greater loss of nucleotide diversity under elevated $p\text{CO}_2$ conditions than in control settings, and the authors suggest that in wild populations, loss of genetic diversity could limit their capacity for further adaptation to future ocean acidification or other drivers in future generations. The authors concluded that whilst some natural populations may currently possess sufficient standing genetic variation to face future global changes, this latent ability of populations to deal with future environmental challenges may be rapidly dissipated by the ongoing environmental change.

The positive effects that trans-generational exposure to stressful conditions appears to exert on a number of marine organisms has led to the idea that human-assisted trans-generational exposure to global change drivers can be used as a conservation tool to produce more resilient offspring (Van Oppen et al 2015). Chakravarti et al. (2016), using an emerging marine polychaete model (*Ophryotrocha labronica*) exposed individuals to projected ocean warming and acidification conditions over successive generations, and showed that trans-generational exposure in the laboratory can produce offspring that are more tolerant when exposed to a single driver; however, trans-generational exposure to combinations of multiple drivers did not always improve traits in the same fashion, potentially due to genetic or physiological constraints or trade-offs. As a consequence, the utilisation of human-assisted acclimation may require an in depth reflection before local and global proactive conservation plans are put into motion (van Oppen et al. 2015).

The existence of trade-offs between tolerance traits to different stressors can limit both species' plastic and evolutionary responses (e.g. Hoffmann and Sgrò 2011; Dam 2013). In order to test this idea, Kelly et al. (2016) hybridized (here intended specifically as crossings) generated from populations of the intertidal copepod *Tigriopus californicus*, differing for both heat and salinity tolerance, followed by a multigenerational selection experiment for tolerance to heat, hypo-osmotic and hyper-osmotic conditions. They found that (i) heat-selected lines were more heat tolerant but showed lower fecundity, (ii) hyper-osmotic-selected lines showed a reduction in tolerance to heat, and (iii), lines selected for both heat and hypo-osmotic stress combined showed a reduced tolerance to heat, this indicating, together with transcriptomic evidence, that energy trade-offs exists for these two tolerance traits.

Finally, in an impressively long-lasting selection experiment, Listmann et al. (2016), investigated changes in thermal reaction norms in the model calcifying coccolithophore *Emiliana huxleyi* in response to 2.5 years of experimental selection to two temperatures (1,200 asexual generations). The different thermal selection regimes led to a marked divergence of thermal reaction norms for optimal growth and maximum persistence temperature to a range of temperatures and $p\text{CO}_2$. Altogether, Listmann et al. (2016)

showed that thermal reaction norms in phytoplankton may evolve at a faster pace than that of predicted ocean warming, bringing some hope for the future of a key element of marine ecosystems.

Epigenetics Responses

Among the mechanisms underlying both plastic and evolutionary processes, especially trans-generational effects, epigenetic mechanisms (e.g., DNA methylation or histone modification) have to date been understudied within the context of marine organisms' responses to global change (Bonduriaski et al. 2012). This may have been primarily caused by the lack of well-developed model organisms and tools for the marine realm, as well as the relatively recent discovery of the importance of these mechanisms. However, many current initiatives address this issue, with new technological advances making it possible to study epigenetic patterns even in less-than-fully developed model systems. Taking advantage of these recent advances, Putnam et al. (2016) tested whether scleractinian corals of the environmentally sensitive species *Pocillopora damicornis* and more environmentally robust species *Montipora capitata* exhibited differences in their phenotypic response that were associated with changes in DNA methylation levels following exposure to elevated $p\text{CO}_2$. Putnam et al. (2016) showed that the more sensitive species exhibited a reduced calcification rate under elevated $p\text{CO}_2$, which was not seen in the more tolerant species. In addition, the sensitive species exhibited larger changes both in its metabolomic profile and DNA methylation pattern, when compared to the most robust species. This novel study highlights the relevance of investigating environmentally induced changes in DNA methylation, as mechanisms mediating the responses to major global change drivers of important ecosystem engineers, such as are corals, whilst at the same time asking relevant broad eco-evolutionary questions. This line of investigation could provide us with a tool to generate heritable plasticity, in support of future conservation actions and to promote assisted evolution in marine organisms (van Oppen et al. 2015). It will be critical to focus future work on the relationship between methylation patterns, gene expression and evolution in the generation of the observed phenotypic trans-generational effects that might provide a rescue mechanism for species facing global change.

The Modelling Approach

Experimental and field observational approaches have so far led the way in building our understanding of how future marine biotas will be shaped by on-going environmental changes (Munday et al. 2013, Godbold and Calosi 2013, Reusch 2013, Sunday et al. 2014). Mathematical models may provide conceptual frameworks within which such experimental data can be placed in context. Further, models can be used as tools in order to design well-

informed and well-designed experiments to produce much needed proof of concepts for key aspects of biological system responses to the global change.

Using an individual based model, Collins (2016) investigated the evolution of cell division rates in asexual populations of unicellular microbes maintained under chronic environmental nutrient enrichment over hundreds of generations. She found that after many generations, initially elevated growth rates appear to become limited by increases in cellular damage. This in turn causes the growth rates to decline to the ancestral state, which Collins (2016) calls the 'Prodigal Son dynamics', in the absence of further evolution for increased tolerance to damages or decreasing in repair cost or decreasing in rate of damages. An implication from this work is that a continuous increase in growth rate, usually taken as a sign of increased fitness, might actually be detrimental to a population in the long run and that intermediate rates are more sustainable and are positively selected for. This theoretical approach is relevant to inform our understanding of how environmental enrichment can increase or control cell division rate in a sustainable fashion, these processes being central to important applications such as biofuel reactors and controlling bio-fouling respectively.

Finally, Marshall et al. (2016) used a heuristic model to explore how traits associated with complex life histories, often found in marine organisms, can alter a population's capacity to cope with environmental change. Marshall et al. (2016) found that an increase in life history complexity decreases the potential for evolution of a species during environmental change. The authors go further, suggesting that levels of genetic correlations in stress tolerance between different life stages, genetic variance levels characterising each life stage, and the relative plasticity level found among different stages, all interact to determine the environmental change threshold any given species can tolerate before extinction occurs. Marshall et al. (2016) concluded based on their model that marine organisms possessing more complex life cycles are particularly sensitive to future global change drivers, but also warn us that for most species we still have to acquire experimental evidence for key traits.

A broader implementation of relatively simple models such as those developed by Collins (2016) and Marshall et al. (2016) could, if well employed and further parameterised with empirical data, rapidly improve our understanding on both specific traits responses and biodiversity responses to the global change.

Conclusion

This special issue collects a number of novel cutting-edge studies showcasing advanced experimental designs, approaches, and tools to be used in the investigation of key aspects of marine organisms' evolutionary responses to ongoing and future rapid global changes. This new knowledge further demonstrates that <<Life may find a way>>, i.e. at least some

populations of some marine species have the ability to adapt to future global change conditions, and illustrates, through transgenerational and epigenetic studies, some of the evolutionary pathways and mechanisms of adaptation that may occur over the next decades. At the same time, we have seen that the potential and capacity of marine organisms for adaptation are finite, due to the presence of evolutionary trade-offs among different traits, particularly when exposed to multiple global change drivers, and the possibility that extant genetic diversity, which enable populations to adapt to changing environments, is quickly reduced with ongoing environmental changes. Consequently, extinction caused by global change in some populations and species in the marine realm, particularly for metazoans, can be expected. This critical understanding of how marine organisms will change under the selective pressure of future global change drivers can be harnessed to help us better predict population, community and ecosystem level responses. This is particularly relevant when consider the discrepancy between our current understanding of the rate of change for environmental parameters *versus* the rate of change (through plasticity and adaptation) of biological systems, within the context of the ongoing global change (Dam 2013, Pörtner et al. 2014). Further studies of species' local adaptation, their capacity for trans- and multigenerational plasticity and rapid evolution, and the existence of epigenetic responses mediating species plasticity and generating diversity upon which selection can act upon, need to be increasingly incorporated into future models of evolution under global change. Such studies will provide powerful tools in our efforts to promote marine conservation, provide increasingly reliable projections on changes in marine biodiversity in the face of global change. At the same time, field observations aiming at detecting ongoing biological changes are critical to assess whether plasticity and adaptation mechanisms are actually occurring in nature, and are occurring at a rate which is fast enough to prevent local and global extinction, also considering the complexity of natural habitats *versus* that of laboratories set ups. This integration will further support our ability to produce reliable projections on changes occurring from the species to the ecosystem level. Current evidence appears to suggest that plasticity and adaptation may not be fully effective in promoting evolutionary rescue (e.g. Pörtner et al., 2014), as past mass extinctions may also suggest, particularly considering the rapidity of the ongoing environmental change (Barnowsky et al. 2011). Nonetheless, relevant evolutionary information will guide us in identifying which conservation efforts may be the most needed to prevent populations and species extinction, and the most effective (i.e. epigenetic manipulation, laboratory transgenerational exposure, artificial selection) (van Oppen et al. 2015). Furthermore this approach will help us identify what rate and magnitude of environmental change we can afford for life to be able to eventually adapt; in turn, which are the thresholds for the rate of the environmental change beyond which evolution will not be effective in rescuing marine organisms? We hope that future efforts (including those by the IPCC) will increasingly incorporate our current, and rapidly increasing, knowledge on marine biological systems' evolutionary responses to rapid environmental changes.

ACKNOWLEDGMENTS

We wish to thank Prof. Hans Dam and Prof. Hans-Otto Pörtner for their constructive criticisms to this manuscript. This work is largely the result of the presentations delivered at the 'Evolutionary Effects of Ocean Warming and Acidification' at the 2015 Aquatic Sciences meeting for the Associating for the Sciences of Limnology and Oceanography, Granada (Spain) 22nd-27th of February 2015, and discussions that took place at the Marine Evolution under Climate Change advance course at The Sven Lovén Centre for Marine Sciences in Kristineberg funded by The Linnaeus Centre for Marine Evolutionary Biology (CeMEB). In addition, this work is a contribution to the work undertaken by CeMEB and the NERC UK Ocean Acidification Research Programme Benthic Consortium task 1.2 and 1.4.

AUTHOR CONTRIBUTIONS

All authors contributed to the rationale for this work. PC, SD and PDW wrote the first draft of this manuscript, and all authors contributed to the final write up.

FUNDING

PC was funded by the NERC UK Ocean Acidification Research Programme (NE/H017127/1), and he is supported by an NSERC Discovery Grant and a FRQ-NT New University Researchers Start Up Program. PDW was supported by a postdoctoral grant from the Marcus and Amalia Wallenberg foundation, as well as BONUS, funded jointly by the EU and the Swedish FORMAS. PT was funded through the "Ocean Acidification and Ecosystem Effects in Northern Waters Flagship" from the High North Research Centre for Climate and the Environment (Fram Centre). SD was financially supported by the Linnaeus Centre for Marine Evolutionary Biology at the University of Gothenburg (<http://www.cemeb.science.gu.se>) and a Linnaeus grant from the Swedish Research Councils VR and Formas.

Reference Lists

1. Barnosky, A. B., Matzke, N., Tomiya, S., Wogan, G. O. U., Swartz, B., Quental, T. B., Marshall, C., McGuire, J. L., Lindsey, E. L., Maguire, K. C., Mersey, B., and E. A. Ferrer 2011. Has the Earth's sixth mass extinction already arrived? *Nature* 471: 51–57.
2. Bennett, A. F., Lenski, R. E., and J. E. Mittler 1992. Evolutionary adaptation to temperature. I. Fitness Responses of *Escherichia coli* to changes in thermal environment. *Evolution* 46, 1: 16-30.
3. Bozinovic, F., Calosi, P., and Spicer, J. I. 2011. Physiological Correlates of Geographic Range in Animals. *Annual Review for Ecology, Evolution and Systematics* 42: 155-179.
4. Bonduriaski, R., Crean, A. J., and T. Day 2012. The implications of nongenetic inheritance for evolution in changing environments. *Evolutionary Applications* 5: 192-201.
5. Calosi, P., Rastrick, S. P. S., Lombardi, C., De Guzman, H. J., Davidson, L., Giangrande, A., Hardege J. D., Schulze, A., Spicer, J. I., Gambi, M.-C. 2013. Adaptation and

acclimatization to ocean acidification in marine ectotherms: an *in situ* transplant with polychaetes at a shallow CO₂ vent system. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 368, 20120444.

6. Chakravarti, L. J., Jarrold, M. D., Gibbin, E. M., Christen, F., Massamba-N'Siala, G., Blier, P. U., and P. Calosi 2016. Can trans-generational experiments be used to enhance species resilience to ocean warming and acidification? *Evolutionary Applications* [Epub ahead of print].
7. Crichton, M. 1990 *Jurassic Park*. Alfred A. Knopf, New York.
8. Colin, S. P., and H. G. Dam 2002. Latitudinal differentiation in the effects of the toxic dinoflagellate *Alexandrium* spp. on the feeding and reproduction of populations of the copepod *Acartia hudsonica*. *Harmful Algae* 1: 113–25.
9. Collins, S. 2016. Growth rate evolution in improved environments under Prodigal Son dynamics. *Evolutionary Applications* [Epub ahead of print].
10. Dam, H. G. 2013. Evolutionary adaptation of marine zooplankton to global change. *Annual Review of Marine Science* 5: 349-370.
11. Darwin, C. 1859. *On the Origin of Species by Natural Selection*. John Murray, London.
12. De Wit, P., S. Dupont, and P. Thor 2015. Selection on oxidative phosphorylation and ribosomal structure as a multigenerational response to ocean acidification in the common copepod *Pseudocalanus acuspes*. *Evolutionary Applications* [Epub ahead of print].
13. Dobzhansky, T. 1937 *Genetics and the Origin of Species*. Columbia University Press, New York.
14. Donelson, J. M., Wong, M., Booth, D. J., and P. L. Munday 2016. Transgenerational plasticity of reproduction depends on rate of warming across generations. *Evolutionary Applications* [Epub ahead of print].
15. Dupont, S., and H.-O. Pörtner 2013. Marine science: Get ready for ocean acidification. *Nature* 498, 7455: 429.
16. Gaston, K. J., Chown, S. L., Calosi, P., Bernardo, J., Bilton, D. T., Clarke, A., Clusella-Trullas, S., Ghalambor, C. K., Konarzewski, M., Peck, L. S., Porter, W. P., Pörtner, H.-O., Rezende, E. L., Schulte, P. M., Spicer, J. I., Stillman, J. H., Terblanche, J. S., and M. van Kleunen 2009. Macrophysiology: A Conceptual Reunification. *American Naturalist* 174: 595-612.
17. Garland, T. and M. R. Rose 1999. *Experimental Evolution: Concepts, Methods, and Applications of Selection Experiments*. University of California Press, Berkeley.
18. Ghalambor, C. K., Mckay, J. K., Carroll, S. P., and D. N. Reznick 2007. Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology* 21, 3: 394–407.
19. Ghalambor, C. K., Hoke, K. L., Ruell, E. W., Fischer, E. K., Reznick, D. N., and K. A. Hughes 2015. Non-adaptive plasticity potentiates rapid adaptive evolution of gene expression in nature. *Nature* 525, 7569: 372-375.

20. Godbold, J., and P. Calosi 2013. Ocean acidification and climate change: advances in ecology and evolution. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 368: 20120448.
21. Gonzalez, A., Ronce, O., Ferriere, R., and M. E. Hochberg 2013. Evolutionary rescue: an emerging focus at the intersection between ecology and evolution. *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 368: 20120404.
22. Hall-Spencer, J. M., Rodolfo-Metalpa, R., Martin, S., Ransome, E., Fine, M., Turner, S. M., Rowley, S. J., Tedesco, D., and M.-C. Buia 2008. Volcanic carbon dioxide vents show ecosystem effects of ocean acidification. *Nature* 454: 96-99.
23. Hoffmann, A. A., and C. M. Sgrò 2011. Climate change and evolutionary adaptation. *Nature* 470: 479–485.
24. Huxley, J. 1942. *Evolution: The Modern Synthesis*. Allen and Unwin, London.
25. IPCC 2013. *Climate Change 2013: The Physical Science Basis: Contribution of Working Group I to the Fifth Assessment Report of the Inter-governmental Panel on Climate Change*. Cambridge University Press, New York.
26. Kellermann, V., van Heerwaarden, B., Sgrò, C. M., and A. A. Hoffmann 2009. Fundamental evolutionary limits in ecological traits drive *Drosophila* species distributions. *Science* 325, 5945: 1244-1246.
27. Kelly, M. W., DeBiase, M. B., Villela, V. A., Roberts, H. L., and C. F. Cecola 2016. Adaptation to climate change: trade-offs among responses to multiple stressors in an intertidal crustacean. *Evolutionary Applications* [Epub ahead of print].
28. Kroeker, K. J., Kordas, R. L., Crim, R., Hendriks, I. E., Ramajo, L., Singh, G. S., Duarte, C. M., and J.-P. Gattuso 2013. Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Global Change Biology* 19: 1884–1896.
29. Lardies M. A., Arias, M. B., Poupin, M. J., Manríquez, P. H., Torres, R., Vargas, C.A., Navarro, J. M., and N. A. Lagos 2014. Differential response to ocean acidification in physiological traits of *Concholepas concholepas* populations. *Journal of Sea Research* 90: 127-134.
30. Listmann, L., LeRoch, M., Schlüter, L., Thomas, M. K., and T. B. H. Reusch 2016. Swift thermal reaction norm evolution in a key marine phytoplankton species. *Evolutionary Applications* [Epub ahead of print].
31. Lohbeck, K. T., Riebesell, U., and T. B. H. Reusch 2012. Adaptive evolution of a key phytoplankton species to ocean acidification. *Nature Geoscience* 5: 346–351.
32. Lloyd, M. M., Makukhov, A. D., and Pespeni, M. H. 2016. Loss of genetic diversity as a consequence of selection in response to high $p\text{CO}_2$. *Evolutionary Applications* [Epub ahead of print].
33. Lucey, N. M., Lombardi, C. DeMarchi, L., Schultze, A., Gambi, M.-C., and P. Calosi 2015. To brood or not to brood: Are marine invertebrates that protect their offspring more resilient to ocean acidification? *Scientific Reports* 5: 12009.

- Accepted Article
34. Lucey, N. M., Lombardi, C., Florio, M., DeMarchi, L., Nannini, M., Rundle, S., Gambi, M. C., and P. Calosi 2016. An *in situ* assessment of local adaptation in a calcifying polychaete from a shallow CO₂ vent system. *Evolutionary Applications* [Epub ahead of print].
 35. Margulis, L. 1999. *Symbiotic Planet: A New Look at Evolution*. Weidenfeld & Nicolson, London.
 36. Marshall, D. J., Burgess, S. C., and T. Connallon 2016. Global change, life-history complexity and the potential for evolutionary rescue. *Evolutionary Applications* [Epub ahead of print].
 37. Melzner, F. Gutowska, M. A., Langenbuch, M., Dupont, S., Lucassen, M., Thorndyke, M. C., Bleich, M., and H.-O. Pörtner 2009. Physiological basis for high CO₂ tolerance in marine ectothermic animals: pre-adaptation through lifestyle and ontogeny? *Biogeosciences* 6, 3: 4693-4738.
 38. Munday, P. L., Warner, R. R., Monroe, K., Pandolfi, J. M., and D. J. Marshall 2013. Predicting evolutionary responses to climate change in the sea. *Evolutionary Applications* 16, 12: 1488–1500.
 39. Noble, D. 2015. Evolution beyond neo-Darwinism: a new conceptual framework. *Journal of Experimental Biology* 218: 7-13.
 40. Padilla-Gamiño, J. L., Gaitán-Espitia, J. D., Kelly, M. W., and G. E. Hofmann 2016. Physiological plasticity and local adaptation to elevated pCO₂ in calcareous algae: An ontogenetic and geographic approach *Evolutionary Applications* [Epub ahead of print].
 41. Pigliucci, M., Murren, C. J., and C. D. Schlichting 2006. Phenotypic plasticity and evolution by genetic assimilation. *Journal of Experimental Biology* 209: 2362-2367.
 42. Pespeni, M. H., Sanford, E., Gaylord, B., Hill, T. M. Hosfelt, J. D., Jarisa, H. K., LaVigner, M., Lenz, E. A., Russell, A. D., Young, M. K., and S. R. Palumbi 2013. Evolutionary change during experimental ocean acidification. *Proceedings of the National Academy of Sciences of the United States of America* 110, 7: 6937–6942.
 43. Pörtner, H.-O., Karl, D. M., Boyd, P. W., Cheung, W., Lluch-Cota, S. E., Nojiri, Y., Schmidt, D. n., Zavialov, P. O., Alheit, J., Aristegui, J., Armstrong, C., Beaugrand, G., Belkovich, V., Bowler, C., Brewer, p., Church, M., Cooley, S. R., del Monte-Luna, P., Edwards, M., Flint, M., Follows, M. J., Frölicher, T., Fulton, E. A., Gattuso, J.-P., Hoegh-Guldberg, O., Hofmann, E. E., Knoll, A. H., Levin, L. A., Menzel, L., Moloney, C. L., Perry, R., Poloczanska, E. S., Roberts, J. M., Rost, B., Sarmiento, J. L., Sedlacek, J., Storch, D., Wiencke, C., and A. C. Wittmann 2016. *Ocean Systems, Chapter 6 Climate Change 2014: impacts, adaptation, and vulnerability. Part A: global and sectorial aspects. Contribution of working group II to the fifth assessment report of the intergovernmental panel on climate change.*
 44. Putnam, H. M., Davidson, J. M., and R. D. Gates 2016 Ocean acidification influences host DNA methylation and phenotypic plasticity in environmentally susceptible corals. *Evolutionary Applications* [Epub ahead of print].

45. Reusch, T.B.H. 2014. Climate change in the oceans: evolutionary *versus* phenotypically plastic responses of marine animals and plants. *Evolutionary Applications* 7: 104-122.
46. Rodriguez-Romero, A., M. D. Jarrold, G. Massamba-N'Siala, J. I. Spicer, and P. Calosi 2015. Multi-generational responses of a marine polychaete to a rapid change in seawater $p\text{CO}_2$. *Evolutionary Applications* [Epub ahead of print].
47. Sanford, E., and M. W. Kelly 2011. Local Adaptation in Marine Invertebrates, *Annual Review of Marine Science* 3: 509-535.
48. Savolainen, O., Lascoux, M., and J. Merilä 2013. Ecological genomics of local adaptation. *Nature Reviews Genetics* 14, 807–820.
49. Shama, L. N. S., F. C. Mark, A. Strobel, A. Lokmer, U. John, and K. M. Wegner 2016. Transgenerational effects persist down the maternal line in marine sticklebacks: gene expression matches physiology in a warming ocean. *Evolutionary Applications* [Epub ahead of print].
50. Somero, G. N. and P. W. Hochachka 2002. *Biochemical Adaptation: Mechanism and Process in Physiological Evolution*. Oxford University Press, New York.
51. Stillman, J. H., and Paganini, A. W. 2015. Biochemical adaptation to ocean acidification. *Journal of Experimental Biology* 218: 1946-1955.
52. Storch, D., Menzel, L., Frickenhaus, S., and H.-O., Pörtner 2014. Climate sensitivity across marine domains of life: limits to evolutionary adaptation shape species interactions. *Global Change Biology* 20, 10: 3059-3067.
53. Sunday, J. M., P. Calosi, S. Dupont, P. L. Munday, J. H. Stillman, and T. B. H. Reusch 2014. Evolution in an acidifying ocean. *Trends in Ecology and Evolution* 29: 117–125.
54. Thor, P., and S. Dupont 2015. Transgenerational effects alleviate severe fecundity loss during ocean acidification in a ubiquitous planktonic copepod. *Global Change Biology* 21: 2261–2271.
55. Van Oppen, M. J. H., J. K. Oliver, H. M. Putnam, and R. D. Gates 2015. Building coral reef resilience through assisted evolution. *Proceedings of the National Academy of Sciences of the United States of America* 112: 2307–2313.
56. Williams, G. C. 1966. *Adaptation and Natural Selection*. Princeton University Press, Princeton.
57. Wittmann, A. C., and H.-O. Poertner 2013. Sensitivities of extant animal taxa to ocean acidification. *Nature Climate Change* 3: 995–1001.
58. Wood, H. L., Sundell, K., Almroth, B. C., Sköld, H. N. and S. P. Eriksson 2016. Population-dependent effects of ocean acidification. *Proceedings of the Royal Society of London B* 283: 20160163.