

Review

Climatic and associated cryospheric, biospheric, and hydrological changes on the Tibetan Plateau: a review

Sadia Bibi,^{a,b} Lei Wang,^{a,b,c}*[®] Xiuping Li,^a Jing Zhou,^a Deliang Chen^{a,d} and Tandong Yao^{a,b,c}

^a Key Laboratory of Tibetan Environmental Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, China

^b University of Chinese Academy of Sciences, Beijing, China
^c CAS Center for Excellence in Tibetan Plateau Earth Sciences, Beijing, China
^d Regional Climate Group, Department of Earth Sciences, University of Gothenburg, Sweden

ABSTRACT: We review recent climate changes over the Tibetan Plateau (TP) and associated responses of cryospheric, biospheric, and hydrological variables. We focused on surface air temperature, precipitation, seasonal snow cover, mountain glaciers, permafrost, freshwater ice cover, lakes, streamflow, and biological system changes. TP is getting warmer and wetter, and air temperature has increased significantly, particularly since the 1980s. Most significant warming trends have occurred in the northern TP. Slight increases in precipitation have occurred over the entire TP with clear spatial variability. Intensification of surface air temperature is associated with variation in precipitation and decreases in snow cover depth, spatial extent, and persistence. Rising surface temperatures have caused recession of glaciers, permafrost thawing, and thickening of the active layers over the permafrost. Changing temperatures, precipitation, and other climate system components have also affected the TP biological system. In addition, elevation-dependent changes in air temperature, wind speed, and summer precipitation have occurred in the TP and its surroundings in the past three decades. Before projecting multifaceted interactions and process responses to future climate change, further quantitative analysis and understanding of the change mechanisms is required.

KEY WORDS climate change; Tibetan Plateau; cryosphere; biosphere; hydrosphere; streamflow

Received 27 June 2017; Revised 14 November 2017; Accepted 15 December 2017

1. Introduction

Tibetan Plateau (TP) is the source of all the major Asian rivers and serves as the main source of water for nearly 1.4 billion people. TP has an average elevation exceeding 4000 m and an area approximately 2.5×10^6 km². TP is covered by snow, glaciers, permafrost, and seasonally frozen ground. It exerts a profound climatic influence on adjacent and distant world regions due to the environmental and climatic influences of the Indian monsoon, East Asian monsoon, and westerlies on the TP (Immerzeel and Bierkens, 2010b; Duan *et al.*, 2012; Yang *et al.*, 2014). In contrast to the Arctic and Antarctic, TP experiences strong interactions among the atmosphere, cryosphere, hydrosphere, and biosphere. As a result, hydrologic and cryosphere changes on the TP caused by climate change directly affect the lives of billions of people.

Climate over the TP is categorized by low air temperatures, high daily temperature differences, low annual temperature difference, and strong solar radiation. A warming rate of 0.46 °C decade⁻¹ was observed during 1984-2009. This is greater than the rate in Northern Hemisphere $(0.38 \degree C \text{ decade}^{-1})$ and almost 1.5 times the rate of global warming $(0.32 \,^{\circ}\text{C} \,\text{decade}^{-1})$ (Zhang *et al.*, 2013a). Recent studies (Liu et al., 2009b; Qin et al., 2006; Pepin, 2015; Guo et al., 2016a, 2016b; Li et al., 2017) have reported that the changing trends in air temperature, wind speed, and summer precipitation are elevation-dependent in the TP and its surroundings. Because of its high sensitivity to climate change, TP changes are considered as an indicator of global warming effects (Yao et al., 2012b). Substantial glacial recession, snow melt, and permafrost degradation have occurred on the TP due to temperature increases which have led to modifications in the hydrological cycle and changes in water resources throughout the region (Jiao et al., 2015).

The period of frozen soil has decreased approximately $15 \text{ day decade}^{-1}$ as a result of permafrost degradation from 1988 to 2007 (Liu *et al.*, 2012). Long-term observations and reanalysis data have been used to evaluate spatio-temporal variations in temperature, precipitation, evapotranspiration (ET), and stream flow and to study TP

^{*} Correspondence to: L. Wang, Key Laboratory of Tibetan Environmental Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, No. 16 Lincui Road, Chaoyang District, Beijing 100101, China. E-mail: wanglei@itpcas.ac.cn

The copyright line for this article was changed on 30 April 2018 after original online publication.

^{© 2018} The Authors. *International Journal of Climatology* published by John Wiley & Sons Ltd on behalf of the Royal Meteorological Society. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.



Figure 1. Responses of hydrosphere, cryosphere, and biosphere to changing climatic system on the TP. SOS means the start of vegetation growing season and EOS represents end of vegetation growing season.

energy and water cycles. Recent studies have focused on different climate variables (Yang et al., 2014; Gao et al., 2015). To develop a regional strategy for climate change and water resource management over TP, it is vital to study hydrological, cryospheric, and biological responses (Chen et al., 2015). Many other reviews have discussed climate change and its impacts on various components of the TP climate system. Kang et al. (2010) reviewed the climate elements that were indicators for cryospheric changes on the TP. Cuo et al. (2014) reviewed the characteristics of surface hydrology and changes in streamflow hydrology over the TP. Yang et al. (2014) studied the impacts of climate change on the TP energy and water cycles. Shen et al. (2015b) studied the link between plant phenology and climate change and quantified the influence of phenological changes on TP ecosystems. Kuang and Jiao (2016) reviewed long-term trends in climate variables during last half century responsible for climate changes on the TP.

In contrast to previous reviews, we provide a comprehensive overview of climate change on TP and its associated hydrological, cryospheric, and biological changes in a systematic and integrated manner. This study evaluates current information and focuses on overall climate change characteristics and responses of the hydrosphere, biosphere, and cryosphere. Most intensive multi-spherical interactions take place on TP, impacting the life of about 20% of the world population. Figure 1 shows how changes in different climatic variables affect biosphere, hydrosphere, and cryosphere and how multi-spheric interactions take place.

2. Changes in the climatic indicators over the TP

2.1. Air temperature

Yearly mean temperature across TP is <0 °C and decreases from east to west (-10 °C in the western TP) (Sun *et al.*,



Figure 2. Trends in annual mean air temperature $(1970-2012, ^{\circ}C \text{ decade}^{-1})$ (a), precipitation $(1970-2012, \text{ mm decade}^{-1})$ (b), wind speed $(1970-2012, \text{ ms}^{-1} \text{ decade}^{-1})$ (c), solar radiation $(1984-2006, \text{ Wm}^{-1} \text{ decade}^{-1})$ (modified from Yang *et al.*, 2011) (d), and relative humidity $(1970-2012, \% \text{ decade}^{-1})$ (e) over the TP.

2015). Surface air temperature from 1970 to 2012 has shown rapid warming (Figure 2(a)). After 1997, the warming trend accelerated (0.25 °C decade⁻¹) compared with an increase during 1980–1997 (0.21 °C decade⁻¹) (Duan and Xiao, 2015). The warming trend of the minimum temperature (0.41 °C decade⁻¹) is around twice the rate of the maximum temperature (0.18 °C decade⁻¹) (You *et al.*, 2016).

Most studies demonstrate that TP air temperature was high in the beginning of 1960s, from mid-1960s to the 1980s it decreased, and from the late 1980s onwards there is sustained increase in air temperature. Model projections have suggested more pronounced increases in minimum temperature than maximum temperatures, and greater winter warming than summer (Liu *et al.*, 2009b). Surface water vapour was identified as one of the main contributors to winter warming on the TP (Rangwala *et al.*, 2009). These will not only accelerate the melting of glaciers that affect the water supply to rivers but downstream precipitation might also be altered (Wang *et al.*, 2008). Nevertheless, the driving mechanisms for future temperature rise have not been revealed completely.

2.2. Precipitation

Similar to air temperature, precipitation is a vital climatic factor in the TP that impacts the cryospheric, biospheric, and hydrological regimes. For instance, the intensity, amount, duration, and phase of precipitation, alongside other characteristics such as its spatial, temporal, and elevational distribution and seasonality, control the development and formation of seasonal snowpack, glacier accretion, watershed storage conditions, soil moisture, and eventually runoff, as hydrological response. This is particularly important for the vulnerable region in the TP under the background of climate warming, which illustrates strong connections between the cryosphere, hydrosphere, and biosphere (Wu et al., 2015; Dong et al., 2016). TP climate is categorized by wet and humid summers with cool and dry winters. Gradual decrease in summer precipitation from southeast to northwest is approximately 700-50 mm. Since the 1960s, the TP has experienced a slight increase in precipitation (Xie et al., 2010; Yang et al., 2011; Gao et al., 2014). This phenomenon is not as pronounced as temperature (Xie et al., 2010). Variation in annual mean precipitation from 1970 to 2012 has been shown in Figure 2(b). The majority of the stations in semi-arid and humid-zones of central TP shows increasing precipitation trends, while at the periphery, decreasing trends have been observed. Summer precipitation increased with elevation (1000-4500 m) over TP from 1970 to 2014 at a rate of 0.83% decade⁻¹ km⁻¹ (Li *et al.*, 2017). The moisture transports from the west by westerlies and from the southwest by the Indian summer monsoon likely contributed the most to the precipitation over the TP (Zhang et al., 2017a). However, complex topography and remoteness of most parts of the TP make it difficult to acquire ground-based meteorological observations.

The increase in atmospheric moisture over TP may be affected by a variety of mechanisms. Xu *et al.* (2008) proposed that water vapour transport increased over TP due to escalated monsoonal circulation and TP warming. Zhang *et al.* (2013a) suggested that melting of glaciers and permafrost due to a warming climate and slight growth in vapour transport from low altitude and Arabian Sea increased the moisture budget over the TP. Other possible mechanisms could involve the polewards shift of the westerly jet and an increase in summer monsoon circulations (Gao *et al.*, 2014).

2.3. Wind speed

Wind speed, as a function of horizontal pressure or temperature gradients, has direct or indirect effects on climatic variations (e.g. sensible heat flux and evaporation) (Guo *et al.*, 2016b). Wind speed has direct or indirect effects on climatic variations, e.g. density of sensible heat flux. Large-scale spatio-temporal variations in global temperature trends, especially the Elevation Dependent Warming (EDW), could be responsible for near-surface wind speed changes (Klink, 1999). The weakening of wind speed is a common phenomenon in China, particularly over TP (Lin et al., 2013) (Figure 2(c)). The annual mean wind speed increases from southeast to northwest. During 1980-2005, differences in annual mean wind speeds were greater in the southeastern TP (1.5 m s^{-1}) than northwestern TP (3.5 m s^{-1}) and the annual entire TP mean value of 2.3 m s⁻¹ was observed (You et al., 2014). Generally, wind speed has decreased on the TP since the 1960s (Kuang and Jiao, 2016) or 1970s (Yang et al., 2011) (Figure 2(c)). Guo et al. (2016b) proposed that the rate of wind speed reduction intensified with elevation with higher-elevation environments having greater changes in wind speed than lower-elevation areas. They suggested that EDW and increased surface roughness may have contributed to the elevation-dependent wind speed reduction in the TP.

2.4. Solar radiation

Worldwide declining trends in solar radiation have occurred at many locations since the 1960s, while pervasive decrease has been seen since the 1990s (Wild *et al.*, 2005). Over the TP, brightening was not observed (Figure 2(d)). The daily mean sunshine duration ranged from 6.8 to 7.4 h over TP (Liang *et al.*, 2013). Since the 1960s, a general decrease in sunshine duration was observed over TP (Yang *et al.*, 2014). From 1961 to 2005 decrease in seasonal mean sunshine durations of -5.7, -7.4, -3.4, and -3.1 h decade⁻¹ was observed during spring, summer, autumn, and winter, respectively, with a maximum decrease in spring and summer (You *et al.*, 2010a, 2010b, 2010c). Since the 1980s, a significant trend of decreasing seasonal sunshine duration has been seen in summer and spring (Kuang and Jiao, 2016).

Changes in solar radiation on the TP have been influenced by anthropogenic aerosol emissions. Increased aerosol emissions have resulted in decreased solar radiation (You et al., 2010a, 2010b, 2010c; Kuang and Jiao, 2016). During the past few decades, the summer monsoon has moved brown clouds from their origin (particularly the southern part of TP) to TP which may have also caused reduction in the amount of solar radiation (Ramanathan et al., 2007; Zhang et al., 2009). You et al., 2010a, 2010b, 2010c found that solar radiation variation is associated with fluctuations of cloud cover over the TP. Yang et al. (2012) proposed that increased amounts of water vapour due to rapid warming and heavy cloud cover are key factors responsible for solar reduction on the TP. Further studies on the effects of decreased solar radiation on the TP are needed.

2.5. Relative humidity

During 1961–2013, mean annual relative humidity has decreased on TP at a rate of -0.23% decade⁻¹ (Yin *et al.*, 2013). However, variability in annual relative humidity

has been observed from the 1960s to 1990s, which has relatively decreased from the 1990s to the present (Kuang and Jiao, 2016).

From the 1960s to the early 21th century the northern TP showed an increasing trend in relative humidity (Zhang, 2007; Yin *et al.*, 2010). Linear trends of relative humidity during 1970–2012 showed large spatial heterogeneity (Figure 2(e)). Spatial distributional trends in relative humidity are complex, and both decreasing trends and increasing trends have occurred throughout the eastern and central part of TP (Kuang and Jiao, 2016). Over the entire region from 1961 to 2013, the seasonal mean relative humidity showed a decreasing trend (apart from winter). The trends were -0.11, -0.60, -0.39, and 0.04% decade⁻¹ in spring, summer, autumn, and winter, respectively (Kuang and Jiao, 2016).

2.6. Causes of climate warming on the TP

The climate warming on the TP may be a result of the interaction of many different factors.

- 1. The recent climate warming on TP is primarily caused by the increased in anthropogenic greenhouse gas emissions (Guo and Wang, 2011; Duan *et al.*, 2012) due to change in local radiative forcing (Duan *et al.*, 2006). Big altitudinal difference and unique topography make the impacts of Green House Gases (GHGs) emissions on TP climate more devastating than any other region (Duan *et al.*, 2006, 2012).
- 2. Snow/ice-albedo and the cloud-radiation feedback are the main contributing factors to climate warming on TP (Rangwala *et al.*, 2010; Duan and Xiao, 2015), where the warming rate in high mountains can be more than in lower-elevation regions. Decreased snow cover due to increased absorbed solar radiation could be part the reasons for the substantial warming over TP.
- 3. Increased ozone concentration could be the possible reasons for decrease in total cloud cover over TP; however, it has been difficult to identify the mechanism (Zhang, 2007). On the other hand, the low-level cloud volumes display antagonistic trend, which decreases during daytime but increases at night (Duan and Wu, 2006). Increased night-time cloud cover could be one of the reasons for increased nocturnal warming in the TP.
- 4. Atmospheric aerosols can affect the atmospheric circulation over TP. Aerosols could increase lower atmospheric solar heating by nearly 50% (Ramanathan *et al.*, 2007; Kang *et al.*, 2010). Aerosols may be transported to the TP by atmospheric circulation (You *et al.*, 2010c) but it is uncertain how these influence the climate warming in the TP. Atmospheric aerosols could modify the regional climate, cryosphere, and hydrology by increased climate warming (Liu *et al.*, 2017). Aerosol particles impact the radiative forcing directly through absorption and reflection of solar and infrared radiation in the atmosphere. Aerosols appear to be more easily transported to the TP across the northern edge rather than the southern edge (Xu *et al.*, 2015).

5. Land use change may also account for climate warming over the TP. Most pronounced land use changes that not only impact the local climate but also have important hydrological implications on TP include grassland and permafrost degradation, deforestation, urbanization, and desertification (Cui and Graf, 2009). Anthropogenic land use change makes the climate warming on TP spatially intricate.

The mechanisms of climate warming on the TP are complex, so at present it is difficult to fully understand. Beside these, surface water vapour and changes in atmospheric circulation and atmospheric aerosol concentration are also considered as important factors that contributed to enhanced climate warming on the TP.

3. Responses of cryosphere to climate change over the TP

3.1. Snow cover

For Asian rivers originating at high altitudes, runoff from snow melt accounts for a greater fraction of the total runoff compared to glacier runoff in Yangtze, Yellow, Mekong, Brahmaputra, and Salween (Zhang et al., 2013b). Precise mapping of the snow cover area is vital for agricultural production and mitigation of snow caused disasters such as floods (Liang et al., 2008). Seasonal snow cover is a key environmental component on the TP and Himalayas regions. Because of cold climatic conditions and high elevation, precipitation usually occurs in solid form. In addition, as a result of low thermal conductivity and high albedo, the snow provides the water and energy interface between the land surface and atmosphere (Shrestha et al., 2012). Snow cover in the southern TP and western edges is most persistent due to Indian monsoon spills onto the plateau (Pu et al., 2007). The monsoonal precipitation (particularly inland) is predicted by spring snow cover on the TP (Immerzeel and Bierkens, 2010a). In contrast to extra tropical areas, snow cover thickness has increased in the TP from 1957 to 1992 (Kang et al., 2010), which coincides with the recent increase in Antarctic and Greenland snow accumulation (Li, 1996).

With greater seasonal variation in spatial extent, snow cover over TP is a distinctive feature, which is a vital source of freshwater for western China and other Himalayan regions. General circulation and monsoon systems over eastern and southern Asia during spring and summer have a strong link with winter snow cover over the TP (Dickson, 1984; Yang, 1996). In regional climate studies, response of snow cover variability in TP to global warming has created an alarming situation (Qin *et al.*, 2006). Most persistent snow cover is situated in western and southeastern edges of TP and it persistency increases with increase in terrain elevation (Pu *et al.*, 2007).

An investigation carried out on snow cover dynamics of four typical lake basins in TP by using MODIS snow cover products (moderate-resolution imaging spectrometer) showed no evident trend of snow cover

S. BIBI et al.



Figure 3. Time series of snow cover variations on the entire TP observed from MOD10A2, IMS, and TAI between 2004 and 2013 (modified from Yu *et al.*, 2016).

change during 2001-2010 (Zhang et al., 2012). Because of biennial variations of the Indian monsoon, in western part of Nam Co Basin (central TP) monsoon and autumn snow cover indicated altering patterns. While snow-free areas and areas with permanent snow cover, snow cover trend was relatively stable (Kropacek et al., 2010). From 1975 to 2000, increased snow depth over the TP was concurrent with a strong Indian-Burma trough, subtropical westerly jet intensification, and increased ascending air mass over the plateau during most of the year (Zhang et al., 2004; Kang et al., 2010). This phenomenon could influence the Asian summer monsoon due to strong snow and monsoon interactions (Ding et al., 2014). As for other climatic factors, it is difficult to understand the duration of snowfall and seasonal snow cover changes on the TP due to limited availability of data on surface snow cover and depth (Li et al., 2008). Yu et al. (2016) compared a time series of snow cover variations crosswise in the TP using MOD10A2, the Interactive Multi-sensor Snow and Ice Mapping System (IMS), and Terra–Aqua–IMS (TAI) between 2001 and 2013. Some MOD10A2 snow cover values (Figure 3) are >40%, while the daily TAI revealed more fine-grained snow cover variations, predominantly higher peaks of snow cover >50%.

3.2. Permafrost

Permafrost covers about 75% of the TP (Cheng and Jin, 2013). Permafrost is a chief component of the cryosphere and susceptible to climate change on different spatial and temporal scales (Cheng and Wu, 2007). In cold regions, thawing depth of frozen soil influences the runoff generation on hillslopes which substantially differs in temperate regions (Zhang *et al.*, 2013c). During the 1970–1990s, an increase in surface temperature over TP from 0.2 to 0.5 °C has caused permafrost reduction and permafrost loss is expected to continue in the future (Zhao *et al.*, 2004;

Lemke *et al.*, 2007). Figure 4 shows the spatial distribution of trends in seasonally frozen ground during 1981–2010 for maximum freezing depths, freeze starting and ending dates, and the freeze duration at a depth of 1 m (Guo and Wang, 2013). Decreasing trends in maximum freezing depth have been observed in TP except for some southern regions. Similarly, freeze start, freeze end, and freeze duration have experienced delaying, advancing, and shortening trends except for some southern parts.

From the 1970s to 1990s, a preliminary assessment was made about of 100 000 km² areal reduction of permafrost in TP (Li *et al.*, 2008). Since the 1970s, permafrost areal extent has decreased about 36% along the Qinghai–Tibetan highway (from Amdo to Liangdaohe) (Lemke *et al.*, 2007). Based on long-term measurements of temperature, the permafrost altitudinal limit has increased 25 m in the northern TP during past 30 years, and increased 50–80 m in southern TP over the past 20 years (Cheng and Wu, 2007). Taliks (i.e. patches of unfrozen ground in an area of permafrost) areal extent stretched approximately 1.2 km alongside the highway. Areal extent of permafrost could shrink more rapidly over the next 50–100 years as predicted by a permafrost numerical model (Nan *et al.*, 2005).

Since the early 1980s, active layer thickness in TP (alongside the Qinghai–Tibetan highway) has increased about 1.0 m from its mean value (Zhao *et al.*, 2004), which is similar in magnitude (0.15–0.50 m) to other places over TP between 1996 and 2001 (Cheng and Wu, 2007). Model results indicate that the active layer thickness in TP has increased about 0.3 m in the north and 0.9 m in locally selected parts during 1980–2001 (Oelke and Zhang, 2007). From 1967 to 1997 thickness of seasonally frozen ground has been reduced by 0.05–0.22 m, and as a result of the earlier onset of spring thaw, duration of seasonally frozen ground has decreased by more than



Figure 4. Trends in the simulated active layer thickness (m decade⁻¹) (a), date of freeze start at 1 m depth (day decade⁻¹) (b), date of freeze end at 1 m depth (day decade⁻¹) (c), and freeze duration at 1 m depth (day decade⁻¹) (d) for permafrost from 1981 to 2010 (modified from Guo and Wang, 2013).

20 days in the TP (Zhao *et al.*, 2004). Along with a climate change indicator, ground freeze-thaw cycles increased the heat-exchange intensity between the ground surface and the atmosphere, which expectedly increased the monsoon on TP (Chang *et al.*, 2007). Thawing can also result in earlier release of trapped carbon dioxide into the atmosphere (Zhang, 2007). Studies on thawing of seasonally frozen ground should be a priority in order to understand possible climatic feedbacks.

3.3. Glaciers

Recent changes in regional climatic conditions have had a negative impact on mass balance due to acceleration of glacier retreat (Kang et al., 2007). By losing 4.5% of combined areal coverage, about 80% of glaciers have receded in western China (Ding et al., 2006). This phenomenon is very evident in the Himalayas (Jin et al., 2005; Bolch et al., 2012; Immerzeel et al., 2012, 2014), Qilian Mountains (Liu et al., 2000; Liu et al., 2002), and Tianshan Mountains (Liu et al., 2006), with a shrinkage rate of between 5 and 10% during last 30 years. In contrast, glaciers in the interior TP are moderately stable (Liu et al., 2000; Liu and Chen, 2000). Nevertheless, accelerated rate of glacier shrinkage has been reported in recent years (Shi, 2001; Li et al., 2008). To present a comprehensive and systematic assessment of glacier status in and around TP, the whole region was divided into seven subregions shown clockwise as I-VII (Figure 5). The Himalaya shows maximum glacial reduction in both length and glacial areal extent (Figures 5(a) and (b)). Like glacial length and areal changes, the most negative mass balances also occur along the Himalayas ranging from -1100 to -760 mm year⁻¹ with an average of $-930 \text{ mm year}^{-1}$ (Figure 5(c)) (Yao et al., 2012a).

Since 1990, most TP glaciers have been rapidly shrinking due to global warming (Yao et al., 2013). Satellite imagery and satellite altimetry have shown great variations in glacial mass and extent over the TP (Guo et al., 2016a). From 2003 to 2009, the rate of glacial shrinkage over the Himalayas was measured at 10-30 mm year-(Gardner et al., 2013), while the volume loss rate of 22 and 45 mm year⁻¹ during 1999–2008 was recorded by a digital elevation model in eastern and western Himalayas, respectively (Gardner et al., 2013). In Khumbu Himal, a 5% decrease in ice coverage was observed during 1962–2005, and a highest volume loss during 1992–2005 (Bolch et al., 2008). In comparison to the mass loss over Himalayas, according to GRACE mission observation during 2003-2010, water mass over TP increased by 7 Gt year⁻¹ as a result increase in water level of many lakes in the central TP (Jacob et al., 2012; Song et al., 2013; Zhang et al., 2013a).

Precipitation change is a key factor contributing to the changes of glacial mass balance and this phenomenon is predominantly important in TP and surroundings. The greatest glacial shrinkage in the Himalayas corresponds to the declining precipitation rate accompanied by the weakening Indian monsoon (Molg et al., 2012, 2014; Yao et al., 2012a; Maussion et al., 2014). Glacial retreat has resulted in hydrological changes, including greater river discharges, elevated lake levels, recurrent glacial lake upsurges, flooding, and increased debris flows from glacial, and water resources. These changes have been the focus of many studies (Lemke et al., 2007; Yao et al., 2007; Li et al., 2008; Zhang et al., 2013a; Cuo et al., 2014; Yang et al., 2014; Jiao et al., 2015). In the 1990s, an increased river runoff (>5.5%) from TP caused by glacier retreat was observed, which is even greater in northwestern China (Yao et al., 2007). Remote sensing data for many



Figure 5. Spatial and temporal patterns of glacier status in the TP and surroundings for glacier retreat in 82 glaciers (a), area reduction in 7090 glaciers (b), and mass balance in 15 glaciers (c) (modified from Yao *et al.*, 2012a, 2012b).

lakes were used to quantify the contribution of glacier melt to rising lake level (Yao *et al.*, 2007). Zhang *et al.* (2017c) investigated that glacier mass loss contributes 13% and ground ice melt due to permafrost degradation contributed 12% in annual changes in lake area, level, and volume during 1970s–2015 in TP. Nevertheless, glacial volume changes and the synchronized environmental implications of glacial recession need additional study.

Despite the data limitations noted, observations have shown clear and systematic patterns of change in climatic regime and cryospheric response over TP. The various lines of evidence are consistent and mutually supportive. Warming has been pervasive, especially during winter and spring and at higher latitudes, while changes in precipitation have been more varied, both regionally and seasonally. Slight increase in precipitation has been observed, especially in the southeastern and southwestern parts of the region, and a decline in the fraction of precipitation falling as snow is very likely associated with rising winter/spring temperatures.

Cryosphere is sensitive to increasing temperature, particularly when precipitation occurs at temperatures near 0 °C. These changes are key drivers for extensive reductions in snow depth, duration, snow cover extent, and freshwater ice cover, predominantly in spring. This has led to an earlier occurrence of the spring freshet across the region. Warmer air temperatures are associated with rising permafrost temperatures, thawing and degradation of permafrost, and increasing glacier melt. Recent declines in winter and annual glacier mass balance at eastern and northeastern TP have been attributed to reduced snow accumulation (Deng *et al.*, 2017), but they may also be due to a shift in precipitation phase from snow to rain in some parts. Anomalously warm conditions led to reduced snow packs and early disappearance of snow cover, also leading to end of summer snow lines retreating off the tops of most glaciers and net negative glacier mass balance.

4. Responses of the biosphere to climate changes over the TP

Vegetation phenology is sensitive to climate changes. Changes in temperature and precipitation and other climate variables could alter the length of growing season. In cold regions, the start of the growth season (SOS) requires adequate temperatures (period preceding SOS that initiates vegetative growth) as a threshold to end ecodormancy (suspension of growth caused by factors outside the meristem but within the plant) during the spring season (Chuine *et al.*, 2003). Warming on TP resulted in a marked advance of SOS during 1982–1999 (Yu *et al.*, 2010; Piao *et al.*, 2011). Because of decreasing photoperiod (daylight)



Figure 6. SOS (a) and EOS (b) in steppe and meadow vegetation of the TP in 1982–2006 (modified from Yu *et al.*, 2010) and spatial distribution of temporal trends (day decade⁻¹) of satellite-derived SOS over 1982–2011 (c) (modified from Shen *et al.*, 2015b). SOS is the start of vegetation growing season, and EOS is the end of vegetation growing season. VIP is variable importance plots and MC is model coefficients. [Correction added on 23 January 2018, after first online publication: This figure has been replaced and its caption has been updated in this current version.]

in the fall, endodormancy (non-growing season) follows ecodormancy and it breaks during dormant period typically at temperature lower than 0 °C during the dormant season (Chuine *et al.*, 2003). Hence, insufficient chilling due to climate warming in late autumn and winter could potentially result in delaying the start of the SOS (Shen *et al.*, 2015a). Delay of SOS in herbaceous plants due to climate warming in autumn and winter has not yet been observed because this temperature is adequate to meet chilling requirements (Shen *et al.*, 2014; Chen *et al.*, 2015). Thus, climate warming delayed the end of vegetation growing season (EOS) on TP due to partial positive inter-annual correlation between growing season temperature and EOS for the alpine steppe and meadow (Liu *et al.*, 2016). Figure 6 demonstrates the spatial distribution of satellite derived SOS temporal trends over TP during 1982–2011 (Shen *et al.*, 2015b) and response of the SOS (a–d) and EOS (e–h) in steppe and meadow vegetation to monthly temperature on TP during 1982–2006 (Yu *et al.*, 2010).

An increase in spring temperatures was observed in TP from 2000 to 2010, and significant decrease in SOS was seen in the western part of the plateau (Shen *et al.*, 2015b). Evidence of advancement in SOS due to pre-season precipitation has been found over the TP, and this phenomenon was more evident in arid parts of the region (Shen *et al.*, 2015a). During 1982–1999 extensive advancement in SOS was caused by the increase in pre-season precipitation

and temperature, while a weakly delayed SOS in northwestern parts of TP might have been due to reduced precipitation (Shen *et al.*, 2015b). The EOS was also delayed on the TP by increases in precipitation during the growing season (Liu *et al.*, 2016). *In situ* observations revealed some positive effects of increasing temperature and precipitation on leaf phenology (Chen *et al.*, 2015). The results suggested that projected increase in temperature and precipitation on the TP would expand the growing season by advancing SOS and postponing EOS (Su *et al.*, 2013). There are several factors that could affect TP phenology. These are listed below:

- 1. Greater increase of warming at night on TP than during the day and the associated smaller diurnal temperature range (Piao *et al.*, 2015).
- 2. Intensity and timing of precipitation, freezing and thawing of soil, evaporation, and snowmelt, degradation of permafrost, increasing evaporation and soil water loss due to climate warming, and extended active layer may have reduced water availability to plants on TP due to the diverse soil properties (Hu *et al.*, 2009; Yang *et al.*, 2009; Zhao *et al.*, 2010; Peng *et al.*, 2013).
- 3. Interactions among different climatic factors also affect the phenology of herbaceous plants (Zhao *et al.*, 2010).
- 4. Photoperiod impacts on phenology over TP have not been adequately assessed (Vitasse and Basler, 2013; Laube *et al.*, 2014).
- No obvious connection has been found between winter warming and spring phenology on TP (Shen *et al.*, 2015b).
- Species composition can be altered either by climatic warming or by anthropogenic activities like overgrazing which ultimately affect the vegetation phenology (Harris, 2010; Niu *et al.*, 2010).

In mutualistic relationships, phenological changes also affect the tropic levels in contagious ecosystems with trophic mismatches in consumer resource dynamics and pollinator host mismatches (Kerby and Post, 2013). TP is a habitat of numerous Tibetan animal species (e.g. Grusnigricollis, Ursusarctospruinosus, and Pantholopshodgsonii), some of which are endangered, so preservation of TP vegetation needs particular attention (Shen et al., 2015b) These changes could ultimately lead to changes in the entire climatic system (Richardson et al., 2013). Both SOS and EOS timing substantially influence the annual carbon budget in the Northern Hemisphere which ultimately causes fluctuations in atmospheric CO₂ concentration (Keenan et al., 2014). Some studies suggest that net carbon uptake will be increasing in spring due to warming driven advancement in leaf unfolding, whereas carbon uptake will either increase or decrease due to delayed leaf fall in autumn (Piao et al., 2008; Keenan et al., 2014). However, more studies are needed to understand the phenomenon of phenological changes on TP and how they affect regional carbon balance and energy cycles (Shen et al., 2015b). Land surface biophysical parameters (sensible heat flux, albedo and trans-boundary layer turbulence) are directly modified by phenological changes which lead to changes in energy and water budgets and thus transform the local climate system (Jeong *et al.*, 2009a, 2009b).

Rapid climate change in the past half century has significantly modified vegetation phenology over TP. Lengthening of growing season is mainly caused by the advance in SOS, and delayed EOS. Plant phenology responds stronger to changes in precipitation patterns also; winter precipitation in the form of rainfall rather than snow requires less heat energy to melt and to warm soil and thus may advance the SOS. Changes in clouds cover and light intensity and other climatic factors also affect SOS and EOS. Some species may green up earlier, and others may senesce later, resulting in stretched growing season at both ends (early SOS and delayed EOS). Along with changes in climate variables, changes in permafrost can also have substantial effects on plateau vegetation phenology and it can also be used as an indirect indicator to monitor the change in permafrost. Further investigations are needed to understand biodiversity change in the alpine region under climate change scenario.

5. Responses of hydrosphere to climate change over TP

5.1. Evapotranspiration

Regional ET change patterns can also vary. Northwestern desert basin has lower ET compared to the southeastern forest cover basin (Kang et al., 2010). Annual mean ET values are 359.7 and 306.6 mm in the upper Yellow and upper Yangtze River basins, respectively (Xue et al., 2013). Linear trends of seasonal mean ET for the upper Yellow River basin, upper Yangtze River basin, Qiangtang Plateau, and Qaidam Basin during 1983–2006 are shown in Figure 7. ET for the four large basins showed increasing trends for all four seasons during past three decades consistent with regional warming (Li et al., 2014). Overall, the inter-annual ET decreased from 2000 to 2010 over 42% of the region, mainly in the northwest of the TP. The dominant climatic factor that controls the long-standing variations of ET in the northwest TP was relative humidity. However, temperature drove ET under moist conditions (Song et al., 2017).

5.2. Water storage changes

Because of its high elevation and glaciers (approximately 1.0×10^5 km², Yao *et al.*, 2012a), snow (approximately 41.9×10^9 m³ year⁻¹ water equivalent, Li *et al.*, 2008), and permafrost (approximately 1.5×10^6 km², Guo and Wang, 2013), TP is very important in global climatic and geodynamical studies (Guo *et al.*, 2016a). In the TP and the Himalayas, direct quantification of mass changes of glaciers, surface water, and groundwater is challenging due to data limitations (Bolch *et al.*, 2011; Kaab *et al.*, 2012). Studies conducted over the TP, Himalayas, and neighbouring areas indicate that water mass changes in the whole region are spatially heterogeneous (Guo *et al.*, 2016a). Water mass in the entire region, particularly in



Figure 7. Linear trends of seasonal mean ET (mm decade⁻¹) for the upper Yellow River basin, upper Yangtze River basin, Qiangtang Plateau, and Qaidam Basin under the period of 1983–2006 (modified from Li *et al.*, 2014). The discharge gauges for the upper Yellow and Yangtze Rivers (Tangnaihai and Zhimenda) are given as solid points.



Figure 8. Spatial distribution of changing rates of total water storage (mm month⁻¹) in the TP from 2003 to 2012 (modified from Jiao et al., 2015).

central and northern parts of the TP, increased between 2003 and 2012, while decreasing in the northern parts of India and Himalayas (Rodell *et al.*, 2009; Guo *et al.*, 2016a). Regularization solution Groupe de Recherches en GeodesieSpatiale (GRGS) investigated the mass changes rate (>10 mm year⁻¹) in the central TP and Qaidam Basin (Guo *et al.*, 2016a). Adjacent to the southern boundary of TP, obvious decrease in total water storage was observed during 2002–2012 (Figure 8). Melting of glacial ice and snow caused the decrease in total water storage along the Yarlung Tsangpo River (Jiao *et al.*, 2015).

Massive tectonic uplifting in the region could recompense the mass loss over the TP and ultimately it makes no substantial contribution to GRACE results (Jacob *et al.*, 2012). Satellite altimetry revealed gradual expansion of water levels of many lakes located in the central parts of the TP (Zhang *et al.*, 2011a; Song *et al.*, 2013). Satellite imagery also discovered many new glacial lakes in the plateau and the number of the new lakes tends to be increasing (Wang *et al.*, 2013). It has provided a positive indication about water mass storage in the form of lakes and increasing soil moisture in the TP (Zhang *et al.*, 2013a). However, where did the lake water or the soil moisture originate? In previous sections, we noted that the entire plateau has experienced climatic warming since the 1980s which has altered the entire TP climatic system. Increases in surface air temperature, decreased solar radiation and humidity, increases in precipitation and melting of glaciers, and decreases in snow cover have all contributed to increased water mass over the TP.

5.3. Lakes

Anthropogenic activities have had minimal impacts on the alpine hydrological environment, so these regions are reasonable indicators of climate change. Understanding the response of alpine lakes to changing climate is crucial for accurate modelling of the hydrological and ecological process, and realistic prediction of seasonal and inter-annual water storage changes (Deniz and Yildiz, 2007; Medina *et al.*, 2008). The TP boasts the greatest concentration of high-altitude inland lakes in the world. With a total area of 41 800 km², it has been estimated that more than 1000 lakes distributed on TP are larger than 1 km² (Ma *et al.*, 2010; Witt *et al.*, 2016). Glacier melt has greatly supported the development of many inland TP lakes because these are fed by glaciers located in their catchments (Yao *et al.*, 2007; Zhu *et al.*, 2010; Günther *et al.*, 2016). Conversely, many other climatic factors such as increased precipitation and decreased evaporation from lake surfaces may have contributed significantly to the expansion of lakes which are not fed by glaciers in their catchment (Liu *et al.*, 2009a).

Rapidly rising temperature is the main driving force behind accelerated glacial retraction and thawing of permafrost which has been the chief source for inland lake expansion on the TP (Yao et al., 2007; Zhu et al., 2010). Since 1969, optical satellite imagery obtained from Landsat and other satellites has been used to study surface extent changes for a small number of lakes (Wang et al., 2007). However, most studies have focused on quantitative detection and analysis of variation in either lake surface area or water level changes which do not provide sufficient information about response of lake water balance to changing climate (Wang et al., 2007; Liu et al., 2009a). Furthermore, due to inaccessibility and remoteness of the areas, the few observation stations are located on Lake Qinghai (Li et al., 2005), Namco Lake (Zhang et al., 2011a), Yamdrok Lake (Li and Wang, 2009), and Zabuye Salt Lake (Qi and Zheng, 2006) which can provide direct hydrological observations such as water level change data. The recent study on TP lakes exhibits an increasing trend of lake expansion from south to north and from west to east (Figure 9) (Lei *et al.*, 2014).

Complex climatic and hydrological environments for specific lakes exhibited variation in the different lake basins (Song et al., 2013). For instance, Lake Qinghai located in northeastern TP, primarily supplemented by precipitation runoff, was shrinking until the 2000s despite of slight increase in precipitation; increase in evaporation rate in the basin was three times higher than precipitation (Zhang et al., 2011b). Thus, increase in water storage cannot be simply explained by a large negative budget caused by 'precipitation-evaporation' in Lake Qinghai (Song et al., 2013). Snow cover and glaciers melt have become chief sources of water for some of the lakes (Zhang et al., 2011b). Similarly, levels of many lakes located in the upper Yellow River basin have shown greater variability in response to instabilities in climate features and water releases into the Yellow River (Song et al., 2013).



Figure 9. Changes in lake area on the interior of TP during the period of 1976-2010 (Lei et al., 2014).

© 2018 The Authors. *International Journal of Climatology* published by John Wiley & Sons Ltd Int. J. Climatol. **38** (Suppl.1): e1–e17 (2018) on behalf of the Royal Meteorological Society.

Likewise, in the early 1970s and mid-1980s climate warming in the TP triggered glacier retraction, which resulted in increased water storage in these lakes (Bolch *et al.*, 2010), but water replenishment forces including rigorous evaporation and infrequent precipitation caused many lakes to shrink or even disappear (Song *et al.*, 2013). Distinct change in lake level during 1997/1998 is thought to be driven by large-scale atmospheric circulation changes in response to climate warming in the region (Zhang *et al.*, 2017b).

In addition to use of satellite images, numerical simulations from a basin-wide hydrological modelling can be effective to quantitatively investigate lake expansion/shrinkage over the TP in detail. This is especially true if *in situ* observations can be used to calibrate and validate the hydrological model. Through the process-based modelling strategy for a specific lake basin (including both water area and its surrounding land area), we are able to determine the key factors that contribute to the historical lake expansion/shrinkage (e.g. Jin *et al.*, 2015). Furthermore, this methodology is promising for prediction of lake changes, with help of the projections of future climatic variables (e.g. precipitation, air temperature, and wind speed).

5.4. Streamflow

Climatic variables such as temperature and precipitation regulate streamflow over the TP. During summer, precipitation is a dominant factor contributing to streamflow over the eastern and southeastern TP due to the strong influence of the South Asian monsoons (Yao et al., 2013). However, spatial pattern understanding is required to quantify the long-term streamflow changes for all the basins on the TP. Streamflow peaks with increases in temperature and precipitation over TP (Yao et al., 2012b). Meltwater is the key contributor to streamflow in the western part and melt increases with increased temperatures (Cuo et al., 2014). In the westerly dominated central part of the TP, both meltwater and groundwater are significant contributors to streamflow during summer. The entire TP displays non-homogeneous spatial and temporal streamflow patterns (Cuo et al., 2014). Variation in components and contributors occurs even for the same river system and streamflow changes from sub-basin to basin and from headwaters to downstream areas (Figure 10). Other factors include prevalent climatic system, environmental and watershed setting, and anthropogenic activities.

Temperature also strongly influences streamflow, and increased temperature positively affects the streamflow if the basin is extensively covered with glaciers. Streamflow response to temperature changes also depends on the forms and spatial distributions of precipitation. In the western TP that is controlled by westerly influences, annual precipitation increases in the range of 20–700 mm from the lowlands to the mountains (Mao *et al.*, 2006; Gao *et al.*, 2010). In some areas, low precipitation generates insufficient streamflow, while in mountainous regions precipitation primarily occurs as snow and ice and contributes to streamflow as meltwater.



Figure 10. Trends of streamflow in the major river basins on the TP (modified from Cuo *et al.*, 2014).

Similar to precipitation and temperature, another important factor that significantly affects the streamflow is actual ET. Studies about actual ET on TP are predominantly based on water balance and potential ET adjusted by available moisture content (Zhang, 2007; Cuo *et al.*, 2013b). Long-term observational actual ET on TP are not available but limited data suggest a general increase in actual ET along with spatial variation which would result in less availability of water for streamflow (Yang *et al.*, 2007; Zhang, 2007; Kang *et al.*, 2010). Larger effects of actual ET on streamflow occur during May–October compared to other months and it is the second most important factor that causes seasonal and annual variation in streamflow besides precipitation (Cuo *et al.*, 2013a).

Large-scale atmospheric systems such as the mid-latitude westerlies, East Asia and Indian monsoons, North Atlantic Oscillation, Arctic Oscillation, El Niño-Southern Oscillation (ENSO), and local circulations all play roles in affecting the weather and climate of TP (Yao *et al.*, 2012b; Cuo *et al.*, 2013b; Gao *et al.*, 2014; Lu *et al.*, 2015). Relating climate system indices and streamflow could possibly reveal the influences of the climate systems on streamflow. This would be helpful for understanding the spatial and temporal changes of streamflow over the TP.

Streamflow is affected by all cryospheric components (Kang *et al.*, 2010). On TP, relationship between seasonal distribution of streamflow and frozen soil is positive (Niu *et al.*, 2011). The degree-day modelling approach has been used to study the contribution of glaciers to seasonal streamflow in various TP basins (Su *et al.*, 2016), but the results of these studies evaluating the quantitative contribution of glaciers are inconsistent (Liu *et al.*, 2009a; Immerzeel *et al.*, 2010; Zhang *et al.*, 2013a). Numerous climate factors affect the future water availability in Asian river basins (Immerzeel and Bierkens, 2012; Lutz *et al.*, 2014). In general, glacier contributions are important

primarily for headwaters or basins with large glacier coverage. The relative importance of snow to stream flow in this region is also controversial. Climate warming affected arid regions more severely than humid part of TP. For the assessment of climate change in the Third Pole and surrounding regions, it is particularly important to quantify the response of hydrosphere as water balance largely depends on climate regimes.

6. Conclusions

The TP has experienced rapid warming and increased precipitation over the last three decades, while solar radiation and wind speed have generally declined. In particular, dry areas have had stronger climate fluctuations than wet regions. The observations described here provided extensive and long-term data for examining climate system change and variability on the TP.

Increase in temperature drives changes in the cryosphere, biosphere, and hydrological cycles of the TP, resulting in seasonal streamflow variation. These changes will lead to changes in biodiversity and water shortages in regions with inadequate water storage capability. Spatial distribution of precipitation trends in the TP is more heterogeneous than temperature and there is significant inter-annual variability. Since the last decade of the 1900s, the rate of glacier retreat has accelerated, so more consideration should be given to subsequent hydrological processes, including greater discharges which can result in glacial lake outburst floods, rising lake levels, increased debris flows, and impacts on water resources in the TP.

Pervasive warming during winter and spring and both temporal and spatial variability in precipitation patterns have been observed over the TP. Decreased winter precipitation and the proportion of precipitation falling as snow may be associated with rising temperatures during winter and spring. These changes have resulted in an extensive reduction in snow cover, snow depth, and duration. Warmer air is the driving force behind rising permafrost temperatures which leads to permafrost thawing and increased glacier melting. Greatest reduction in glacial length and area and the most negative mass balance on the TP have been attributed to reduced snow accumulation (Yao et al., 2012a). Because of the few meteorological stations in the western TP, limited solid precipitation data are available for this region. The responses of permafrost thawing and the reduction of seasonally frozen ground in response to changing climate on TP and its hydrological response across watersheds and gradients in different physiographic and thermal environments need to be studied. The influence of declining snowfall and duration on all facets of the hydrological cycle and streamflow will also be useful study topics.

Plant phenology has a broad influence on the structure and function of ecosystems but no model has been applied to study phenological responses on the TP. Because of our limited understanding of biosphere responses to climate change, additional modelling should be carried out to address phenological dynamics and responses to a changing climate such as future water availability. Ongoing research should continue evaluating the interactions between climate, hydrology, ecology, cryosphere, and other components of TP environments.

Acknowledgements

This work was funded by the "Strategic Priority Research Program" of the Chinese Academy of Sciences (XDA19070301) and the National Natural Science Foundation of China (Grant 91747201, 41571033). It was also supported by the International Partnership Program of the Chinese Academy of Sciences (131C11KYSB20160061) and the Top-Notch Young Talents Program of China (Dr. Lei Wang). The authors appreciate Dr. Xiaotao Zhang for providing figure 8.

References

- Bolch T, Buchroithner M, Pieczonka T, Kunert A. 2008. Planimetric and volumetric glacier changes in the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data. J. Glaciol. 54: 592–600.
- Bolch T, Yao T, Kang S, Buchroithner MF, Scherer D, Maussion F, Schneider C. 2010. A glacier inventory for the western Nyainqentanglha Range and the Nam Co Basin, Tibet, and glacier changes 1976–2009. *Cryosphere* **4**: 419–433.
- Bolch T, Pieczonka T, Benn D. 2011. Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery. *Cryosphere* 5: 349–358.
- Bolch T, Kulkarni A, Kaab A, Huggel C, Paul F, Cogley JG, Bajracharya S. 2012. The state and fate of Himalayan glaciers. *Science* **336**: 310–314.
- Chang G, Li L, Zhu X, Wang Z, Xiao J, Li F. 2007. Changes and influencing factors of surface water resources in the source region of the Yellow River. Acta Geogr. Sin. 62(3): 311–320.
- Chen X, An S, Inouye DW, Schwartz MD. 2015. Temperature and snowfall trigger alpine vegetation green up on the world's roof. *Global Changes Biol.* **21**: 3635–3646.
- Cheng G, Jin H. 2013. Permafrost and groundwater on the Qinghai–Tibet Plateau and in northeast China. *Hydrgeol. J.* 21: 5–23.
- Cheng G, Wu T. 2007. Responses of permafrost to climate change and their environmental significance, Qinghai–Tibet Plateau. J. Geophys. Res. Earth 112: F02S03.
- Chuine I, Kramer K, Hänninen H. 2003. Plant development models. In *Phenology: An Integrative Environmental Science*. Springer: Dordrecht, 217–235. https://doi.org/10.1007/978-94-007-0632-3_14.
- Cui X, Graf HF. 2009. Recent land cover changes on the Tibetan Plateau: a review. *Clim. Change* **94**: 47–61.
- Cuo L, Zhang Y, Gao Y, Hao Z, Cairang L. 2013a. The impacts of climate change and land cover/use transition on the hydrology in the upper Yellow River basin, China. *J. Hydrol.* **502**: 37–52.
- Cuo L, Zhang Y, Wang Q, Zhang L, Zhou B, Hao Z, Su F. 2013b. Climate change on the northern Tibetan Plateau during 1957–2009: spatial patterns and possible mechanisms. J. Climatol. 26: 85–109.
- Cuo L, Zhang Y, Zhu F, Liang L. 2014. Characteristics and changes of streamflow on the Tibetan Plateau: a review. J. Hydrol. Reg. Stud. 2: 49–68.
- Deng H, Pepin NC, Chen Y. 2017. Changes of snowfall under warming in the Tibetan Plateau. J. Geophys. Res. Atmos. 122: 7323-7341. https://doi.org/10.1002/2017JD026524.
- Deniz O, Yildiz M. 2007. The ecological consequences of level changes in Lake Van. Water Resour. 34: 707–711.
- Dickson PR. 1984. Eurasian snow cover versus Indian monsoon rainfall an extension of the Hahn-Shukla results. *J. Appl. Meteorol.* 23: 171–173.
- Ding Y, Liu S, Li J, Shangguan D. 2006. The retreat of glaciers in response to recent climate warming in western China. *Ann. Glaciol.* 43: 97–105.
- Ding Y, Liu Y, Liang S, Ma X, Zhang Y, Si D, Zhang J. 2014. Interdecadal variability of the East Asian winter monsoon and its possible links to global climate change. *J. Meteorol. Res.* 28: 693–713.

- Dong W, Lin Y, Wright JS, Ming Y, Xie Y, Wang B, Tian L. 2016. Summer rainfall over the southwestern Tibetan Plateau controlled by deep convection over the Indian subcontinent. *Nat. Commun.* 7: 10925.
- Duan A, Wu G. 2006. Change of cloud amount and the climate warming on the Tibetan Plateau. *Geophys. Res. Lett.* **33**: L22704.
- Duan A, Wu G, Zhang Q, Liu Y. 2006. New proofs of the recent climate warming over the Tibetan Plateau as a result of the increasing greenhouse gases emissions. *China Sci. Bull.* **51**: 1396–1400.
- Duan A, Wu G, Liu Y, Ma Y, Zhao P. 2012. Weather and climate effects of the Tibetan Plateau. Adv. Atmos. Sci. 29: 978–992.
- Duan A, Xiao Z. 2015. Does the climate warming haitus exist over the Tibetan Plateau? *Sci. Rep.* 5: 13711. https://doi.org/10.1038/ srep13711.
- Gao X, Ye B, Zhang S, Qiao C, Zhang X. 2010. Glacier runoff variation and its influence on river runoff during 1961–2006 in the Tarim River basin, China. Sci. China Earth Sci. 53: 880–891.
- Gao Y, Cuo L, Zhang Y. 2014. Changes in moisture flux over the Tibetan Plateau during 1979–2011 and possible mechanisms. J. Climatol. 27: 1876–1893.
- Gao Y, Li X, Leung LR, Chen D, Xu J. 2015. Aridity changes in the Tibetan Plateau in a warming climate. *Environ. Res. Lett.* 10: 034013.
- Gardner AS, Moholdt G, Cogley JG, Wouters B, Arendt AA, Wahr J, Ligtenberg SR. 2013. A reconciled estimate of glacier contributions to sea level rise: 2003 to 2009. *Science* **340**: 852–857.
- Günther F, Thiele A, Biskop S, Mäusbacher R, Haberzettl T, Yao T, Gleixner G. 2016. Late quaternary hydrological changes at Tangra Yumco, Tibetan Plateau: a compound-specific isotope-based quantification of lake level changes. J. Paleolimnol. 55: 369–382.
- Guo D, Wang H. 2011. The significant climate warming in the northern Tibetan Plateau and its possible causes. *Int. J. Climatol.* 32: 1775–1781.
- Guo D, Wang H. 2013. Simulation of permafrost and seasonally frozen ground conditions on the Tibetan Plateau, 1981–2010. J. Geophys. Res. Atmos. 118: 5216–5230.
- Guo J, Mu D, Liu X, Yan H, Sun Z, Guo B. 2016a. Water storage changes over the Tibetan Plateau revealed by GRACE mission. *Acta Geophys.* 64: 463–476.
- Guo X, Wang L, Tian L, Li X. 2016b. Elevation of dependent reductions in wind speed over and around the Tibetan Plateau. *Int. J. Climatol.* 37: 1117–1126. https://doi.org/10.1002/joc.4727.
- Harris RB. 2010. Rangeland degradation on the Qinghai–Tibetan Plateau: a review of the evidence of its magnitude and causes. J. Arid Environ. 74: 1–12.
- Hu H, Wang G, Liu G, Li T, Ren D, Wang Y, Wang J. 2009. Influences of alpine ecosystem degradation on soil temperature in the freezing-thawing process on Qinghai–Tibet Plateau. *Environ. Geol.* 57: 1391–1397.
- Immerzeel W, Bierkens M. 2010a. Seasonal prediction of monsoon rainfall in three Asian river basins: the importance of snow cover on the Tibetan Plateau. *Int. J. Climatol.* **30**: 1835–1842.
- Immerzeel WW, Bierkens MF. 2010b. Asian water towers: more on monsoons response. *Science* 330: 585–585.
- Immerzeel W, Bierkens M. 2012. Asia's water balance. Nat. Geosci. 5: 841–842.
- Immerzeel WW, Van Beek LP, Bierkens MF. 2010. Climate change will affect the Asian water towers. *Science* 328: 1382–1385.
- Immerzeel W, Pellicciotti F, Bierkens M. 2012. Rising river flows throughout the 21st century in two Himalayan glacierised watersheds. *Nat. Geosci.* 6: 742–745.
- Immerzeel W, Kraaijenbrink P, Shea J, Shrestha A, Pellicciotti F, Bierkens M, De Jong S. 2014. High-resolution monitoring of Himalayan glacier dynamics using unmanned aerial vehicles. *Remote Sens. Environ.* 150: 93–103.
- Jacob T, Wahr J, Pfeffer WT, Swenson S. 2012. Recent contributions of glaciers and ice caps to sea level rise. *Nature* 482: 514–518.
- Jeong SJ, Ho CH, Jeong JH. 2009a. Increase in vegetation greenness and decrease in springtime warming over East Asia. *Geophys. Res. Lett.* 36: L02710.
- Jeong SJ, Ho CH, Kim KY, Jeong JH. 2009b. Reduction of spring warming over East Asia associated with vegetation feedback. *Geophys. Res. Lett.* 36: L18705.
- Jiao JJ, Zhang X, Liu Y, Kuang X. 2015. Increased water storage in the Qaidam Basin, the North Tibet Plateau from GRACE gravity data. *Plos One* 10: e0141442.
- Jin R, Li X, Che T, Wu L, Mool P. 2005. Glacier area changes in the Pumqu River basin, Tibetan Plateau, between the 1970s and 2001. *J. Glaciol.* **51**: 607–610.

- Jin Z, An Z, Yu J, Li F, Zhang F. 2015. Lake Qinghai sediment geochemistry linked to hydroclimate variability since the last glacial. *Ouat. Sci. Rev.* 122: 63–73.
- Kaab A, Berthier E, Nuth C, Gardelle J, Arnaud Y. 2012. Contrasting patterns of early twenty-first-century glacier mass change in the Himalayas. *Nature* 488: 495–498.
- Kang S, Qin D, Ren J, Zhang Y, Kaspari S, Mayewski PA, Hou S. 2007. Annual accumulation in the Mt. Nyainqentanglha ice core, southern Tibetan Plateau, China: relationships to atmospheric circulation over Asia. Arct. Antarct. Alp. Res. 39: 663–670.
- Kang S, Xu Y, You Q, Flügel W-A, Pepin N, Yao T. 2010. Review of climate and cryospheric change in the Tibetan Plateau. *Environ. Res. Lett.* 5: 015101.
- Keenan TF, Gray J, Friedl MA, Toomey M, Bohrer G, Hollinger DY, Yang B. 2014. Net carbon uptake has increased through warming-induced changes in temperate forest phenology. *Nat. Clim. Change* 4: 598–604.
- Kerby JT, Post E. 2013. Advancing plant phenology and reduced herbivore production in a terrestrial system associated with sea ice decline. *Nat. Commun.* 4: 2514.
- Klink K. 1999. Trends in mean monthly maximum and minimum surface wind speeds in the coterminous United States, 1961 to 1990. *Clim. Res.* 13: 193–205.
- Kropacek J, Feng C, Alle M, Kang S, Hochschild V. 2010. Temporal and spatial aspects of snow distribution in the Nam Co Basin on the Tibetan Plateau from MODIS data. *Remote Sens.* 2: 2700–2712.
- Kuang X, Jiao JJ. 2016. Review on climate change on the Tibetan Plateau during the last half century. J. Geophys. Res. Atmos. 121: 3979–4007.
- Laube J, Sparks TH, Estrella N, Höfler J, Ankerst DP, Menzel A. 2014. Chilling outweighs photoperiod in preventing precocious spring development. *Global Change Biol.* 20: 170–182.
- Lei Y, Yang K, Wang B, Sheng Y, Bird BW, Zhang G, Tian L. 2014. Response of inland lake dynamics over the Tibetan Plateau to climate change. *Clim. Change* **125**: 281–290.
- Lemke P, Ren J, Alley RB, Allison I, Carrasco J, Flato G, Zhang T. 2007. In Observations: Changes in Snow, Ice and Frozen Ground, Climate Change: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Solomon S, Qin D, Manning M, Chen Z, Marquis M, Averyt KB, Tignor M, Miller HL (eds). Cambridge University Press: Cambridge, UK, 337–383.
- Li P. 1996. Response of Tibetan snow cover to global warming. Acta Geophys. Sin. 51: 260–265.
- Li L, Wang W. 2009. The response of lake change to climate fluctuation in north Qinghai–Tibet Plateau in last 30 years. J. Geogr. Sci. 19: 131–142.
- Li L, Zhu X, Wang Z, Wang Q. 2005. Influential factor of water level variation of Qinghai Lake in recent 42 years and trend prediction. J. Desert Res. 25: 690–696.
- Li X, Cheng G, Jin H, Kang E, Che T, Jin R, Shen Y. 2008. Cryospheric change in China. *Global Planet. Change* 62: 210–218.
- Li X, Wang L, Chen D, Yang K, Wang A. 2014. Seasonal evapotranspiration changes (1983–2006) of four large basins on the Tibetan Plateau. J. Geophys. Res. Atmos. 119: 13079–13095.
- Li X, Wang L, Guo X, Chen D. 2017. Does summer precipitation trend over and around the Tibetan Plateau depend on elevation? *Int. J. Climatol.* 37: 1278–1284. https://doi.org/10.1002/joc.4978.
- Liang TG, Huang XD, Wu CX, Liu XY, Li WL, Guo ZG, Ren JZ. 2008. An application of MODIS data to snow cover monitoring in a pastoral area: a case study in northern Xinjiang, China. *Remote Sens. Environ.* 112: 1514–1526.
- Liang L, Li L, Liu C, Cuo L. 2013. Climate change in the Tibetan Plateau three rivers source region: 1960–2009. *Int. J. Climatol.* 33: 2900–2916.
- Lin C, Yang K, Qin J, Fu R. 2013. Observed coherent trends of surface and upper-air wind speed over China since 1960. J. Climatol. 26: 2891–2903.
- Liu X, Chen B. 2000. Climatic warming in the Tibetan Plateau during recent decades. Int. J. Climatol. 20: 1729–1742.
- Liu C, Shi Y, Wang Z, Xie Z. 2000. Glacier resources and their distributive characteristics in China: a review on Chinese glacier inventory. *J. Glaciol. Geocryol.* **22**: 106–112.
- Liu S, Shen Y, Sun W, Li G. 2002. Glacier variation since the maximum of the Little Ice Age in the western Qilian Mountains, northwest China. *J. Glaciol. Geocryol.* **24**: 227–233.
- Liu SY, Ding YJ, Zhang Y, Shangguan DH, Li J, Han HD, Xie CW. 2006. Impact of the glacial change on water resources in the Tarim River basin. *Acta Oceanol. Sin.* **61**: 482–490.

© 2018 The Authors. *International Journal of Climatology* published by John Wiley & Sons Ltd on behalf of the Royal Meteorological Society.

- Liu J, Wang S, Yu S, Yang D, Zhang L. 2009a. Climate warming and growth of high-elevation inland lakes on the Tibetan Plateau. *Global Planet. Change.* 67: 209–217.
- Liu X, Cheng Z, Yan L, Yin Z. 2009b. Elevation dependency of recent and future minimum surface air temperature trends in the Tibetan Plateau and its surroundings. *Global Planet. Change*. 68(3): 164–174.
- Liu Y, Wu G, Hong J, Dong B, Duan A, Bao Q, Zhou L. 2012. Revisiting Asian monsoon formation and change associated with Tibetan Plateau forcing: II. Change. *Climatol. Dyn.* **39**: 1183–1195.
- Liu Q, Fu YH, Zeng Z, Huang M, Li X, Piao S. 2016. Temperature, precipitation, and insolation effects on autumn vegetation phenology in temperate China. *Global Change Biol.* 22: 644–655.
- Liu B, Cong Z, Wang Y, Xin J, Wan X, Pan Y, Liu Z, Wang Y, Zhang G, Wang Z, Wang Y, Kang S. 2017. Background aerosol over the Himalayas and Tibetan Plateau: observed characteristics of aerosol mass loading. *Atmos. Chem. Phys.* 17: 449–463.
- Lu N, Trenberth KE, Qin J, Yang K, Yao L. 2015. Detecting long-term trends in precipitable water over the Tibetan Plateau by synthesis of station and MODIS observations. *J. Climatol.* **28**: 1707–1722.
- Lutz A, Immerzeel W, Shrestha A, Bierkens M. 2014. Consistent increase in High Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Change* 4: 587–592.
- Ma R, Duan H, Hu C, Feng X, Li A, Ju W, Yang G. 2010. A half-century of changes in China's lakes: global warming or human influence? *Geophys. Res. Lett.* 37: L24106.
- Mao W, Sun B, Wang T, Luo G, Zhang C, Huo L. 2006. Change trends of temperature, precipitation and runoff volume in the Kaxgar River basin since recent 50 years. *Arid Zone Res.* 23: 531–538.
- Maussion F, Scherer D, Mölg T, Collier E, Curio J, Finkelnburg R. 2014. Precipitation seasonality and variability over the Tibetan Plateau as resolved by the High Asia reanalysis. *J. Clim.* **27**: 1910–1927.
- Medina CE, Gomez-Enri J, Alonso JJ, Villares P. 2008. Water level fluctuations derived from ENVISAT radar altimeter (RA-2) and in-situ measurements in a subtropical waterbody: Lake Izabal (Guatemala). *Remote Sens. Environ.* **112**: 3604–3617.
- Molg T, Maussion F, Yang W, Scherer D. 2012. The footprint of Asian monsoon dynamics in the mass and energy balance of a Tibetan glacier. *Cryosphere* **6**: 1445–1461.
- Molg T, Maussion F, Scherer D. 2014. Mid-latitude westerlies as a driver of glacier variability in monsoonal High Asia. *Nat. Clim. Change* 4: 68–73.
- Nan Z, Li S, Cheng G. 2005. Prediction of permafrost distribution on the Qinghai–Tibet Plateau in the next 50 and 100 years. *Sci. China Ser. D Earth Sci.* 48: 797–804.
- Niu K, Zhang S, Zhao B, Du G. 2010. Linking grazing response of species abundance to functional traits in the Tibetan alpine meadow. *Plant Soil* 330: 215–223.
- Niu L, Ye B, Li J, Sheng Y. 2011. Effect of permafrost degradation on hydrological processes in typical basins with various permafrost coverage in western China. *Sci. China Earth Sci.* 54: 615–624.
- Oelke C, Zhang T. 2007. Modeling the active-layer depth over the Tibetan Plateau. Arct. Antarct. Alp. Res. **39**: 714–722.
- Peng S, Piao S, Ciais P, Myneni RB, Chen A, Chevallier F, Vicca S. 2013. Asymmetric effects of daytime and night-time warming on Northern Hemisphere vegetation. *Nature* 501: 88–92.
- Pepin NC. 2015. Elevation-dependent warming in mountain regions of the world. *Nat. Clim. Change* **5**: 424–430.
- Piao S, Ciais P, Friedlingstein P, Peylin P, Reichstein M, Luyssaert S, Grelle A. 2008. Net carbon dioxide losses of northern ecosystems in response to autumn warming. *Nature* 451: 49–52.
- Piao S, Cui M, Chen A, Wang X, Ciais P, Liu J, Tang Y. 2011. Altitude and temperature dependence of change in the spring vegetation green-up date from 1982 to 2006 in the Qinghai–Xizang Plateau. *Agric. For. Meteorol.* 151: 1599–1608.
- Piao S, Tan J, Chen A, Fu YH, Ciais P, Liu Q, Li Y. 2015. Leaf onset in the Northern Hemisphere triggered by daytime temperature. *Nat. Commun.* 6: 6911.
- Pu Z, Xu L, Salomonson VV. 2007. MODIS/Terra observed seasonal variations of snow cover over the Tibetan Plateau. *Geophys. Res. Lett.* 34: L06706.
- Qi W, Zheng M. 2006. Initial research on water level fluctuation discipline of Zabuye Salt Lake in Tibet. *Geogr. Sci.* 26: 693–701.
- Qin D, Liu S, Li P. 2006. Snow cover distribution, variability and response to climate change in western China. J. Clim. 19: 1820–1833.
- Ramanathan V, Ramana MV, Roberts G, Kim D, Corrigan C, Chung C, Winker D. 2007. Warming trends in Asia amplified by brown cloud solar absorption. *Nature* 448: 575–578.

- Rangwala I, Miller JR, Xu M. 2009. Warming in the Tibetan Plateau: possible influences of the changes in surface water vapor. *Geophys. Res. Lett.* 36: L06703.
- Rangwala I, Miller JR, Russell GL, Xu M. 2010. Using a global climate model to evaluate the influences of water vapor, snow cover and atmospheric aerosol on warming in the Tibetan Plateau during the twenty-first century. *Clim. Dyn.* 34: 859–872.
- Richardson AD, Keenan TF, Migliavacca M, Ryu Y, Sonnentag O, Toomey M. 2013. Climate change, phenology, and phenological control of vegetation feedbacks to the climate system. *Agric. For. Meteorol.* **169**: 156–173.
- Rodell M, Velicogna I, Famiglietti JS. 2009. Satellite-based estimates of groundwater depletion in India. *Nature* 460: 999–1002.
- Shen M, Zhang G, Cong N, Wang S, Kong W, Piao S. 2014. Increasing altitudinal gradient of spring vegetation phenology during the last decade on the Qinghai–Tibetan Plateau. Agric. For. Meteorol. 189: 71–80.
- Shen M, Piao S, Cong N, Zhang G, Jassens IA. 2015a. Precipitation impacts on vegetation spring phenology on the Tibetan Plateau. *Global Change Biol.* 21: 3647–3656.
- Shen M, Piao S, Dorji T, Liu Q, Cong N, Chen X, Zhang G. 2015b. Plant phenological responses to climate change on the Tibetan Plateau: research status and challenges. *Natl. Sci. Rev.* **2**: 454–467.
- Shi Y. 2001. Estimation of the water resources affected by climatic warming and glacier shrinkage before 2050 in west China. J. Glaciol. Geocryol. 23: 333–341.
- Shrestha M, Wang L, Koike T, Xue Y, Hirabayashi Y. 2012. Modeling the spatial distribution of snow cover in the Dudhkoshi region of the Nepal Himalayas. J. Hydrometeorol. 13: 204–222.
- Song C, Huang B, Ke L. 2013. Modeling and analysis of lake water storage changes on the Tibetan Plateau using multi-mission satellite data. *Remote Sens. Environ.* 135: 25–35.
- Song L, Zhuang Q, Yin Y, Zhu X, Wu S. 2017. Spatio-temporal dynamics of evapotranspiration on the Tibetan Plateau from 2000 to 2010. *Environ. Res. Lett.* 12: 014011.
- Su F, Duan X, Chen D, Hao Z, Cuo L. 2013. Evaluation of the global climate models in the CMIP5 over the Tibetan Plateau. J. Climatol. 26: 3187–3208.
- Su F, Zhang L, Ou T, Chen D, Yao T, Tong K, Qi Y. 2016. Hydrological response to future climate changes for the major upstream river basins in the Tibetan Plateau. *Global Planet. Change* **136**: 82–95.
- Sun Q, Miao C, Duan Q. 2015. Projected changes in temperature and precipitation in ten river basins over China in 21st century. *Int. J. Climatol.* 35: 1125–1141.
- Vitasse Y, Basler D. 2013. What role for photoperiod in the bud burst phenology of European beech. Eur. J. For. Res. 132: 1–8.
- Wang L, Lu A, Yao T, Wang N. 2007. The study of typical glaciers and lakes fluctuations using remote sensing in Qinghai–Tibetan Plateau. In *IEEE International Geoscience and Remote Sensing Symposium*, 23–28 July 2007, Barcelona, Spain, 4526–4529.
- Wang B, Bao Q, Hoskins B, Wu G, Liu Y. 2008. Tibetan Plateau warming and precipitation changes in East Asia. *Geophys. Res. Lett.* 35: L14702.
- Wang X, Siegert F, Zhou A-G, Franke J. 2013. Glacier and glacial lake changes and their relationship in the context of climate change, central Tibetan Plateau 1972–2010. *Global Planet. Change* **111**: 246–257.
- Wild M, Gilgen H, Roesch A, Ohmura A, Long CN, Dutton EG, Tsvetkov A. 2005. From dimming to brightening: decadal changes in solar radiation at Earth's surface. *Science* **308**: 847–850.
- Witt R, Günther F, Lauterbach S, Kasper T, Mäusbacher R, Yao T, Gleixner G. 2016. Biogeochemical evidence for freshwater periods during the last glacial maximum recorded in lake sediments from Nam Co, south-central Tibetan Plateau. J. Paleolimnol. 55: 67–82.
- Wu G, Duan A, Liu Y, Mao J, Ren R, Bao Q, Hu W. 2015. Tibetan Plateau climate dynamics: recent research progress and outlook. *Natl. Sci. Rev.* 2: 100–116.
- Xie H, Ye J, Liu X, Chongyi E. 2010. Warming and drying trends on the Tibetan Plateau (1971–2005). *Theor. Appl. Climatol.* **101**: 241–253.
- Xu X, Lu C, Shi X, Gao S. 2008. World water tower: an atmospheric perspective. *Geophys. Res. Lett.* **35**: L20815.
- Xu C, Ma YM, You C, Zhu ZK. 2015. The regional distribution characteristics of aerosol optical depth over the Tibetan Plateau. *Atmos. Chem. Phys.* 15: 12065–12078.
- Xue B, Wang L, Li X, Yang K, Chen D, Sun L. 2013. Evaluation of evapotranspiration estimates for two river basins on the Tibetan Plateau by a water balance method. J. Hydrol. 492: 290–297.
- Yang S. 1996. ENSO-snow-monsoon associations and seasonal-interannual predictions. Int. J. Climatol. 16: 2038–2051.

- Yang J, Ding Y, Chen R. 2007. Climatic causes of ecological and environmental variations in the source regions of the Yangtze and Yellow Rivers of China. *Environ. Geol.* **53**: 113–121.
- Yang K, Chen Y, Qin J. 2009. Some practical notes on the land surface modeling in the Tibetan Plateau. *Hydrol. Earth Syst. Sci.* 13: 687–701.
- Yang K, Ye B, Zhou D, Wu B, Foken T, Qin J, Zhou Z. 2011. Response of hydrological cycle to recent climate changes in the Tibetan Plateau. *Clim. Change* 109: 517–534.
- Yang K, Ding B, Qin J, Tang W, Lu N, Lin C. 2012. Can aerosol loading explain the solar dimming over the Tibetan Plateau? *Geophys. Res. Lett.* 39: L20710.
- Yang K, Wu H, Qin J, Lin C, Tang W, Chen Y. 2014. Recent climate changes over the Tibetan Plateau and their impacts on energy and water cycle: a review. *Global Planet. Change* **112**: 79–91.
- Yao T, Pu J, Lu A, Wang Y, Yu W. 2007. Recent glacial retreat and its impact on hydrological processes on the Tibetan Plateau, China, and surrounding regions. Arct. Antarct. Alp. Res. 39: 642–650.
- Yao T, Thompson L, Yang W, Yu W, Gao Y, Guo X, Pu J. 2012a. Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings. *Nat. Clim. Change* 2: 663–667.
- Yao T, Thompson LG, Mosbrugger V, Zhang F, Ma Y, Luo T, Joswiak ME. 2012b. Third pole environment (TPE). *Environ. Dev.* 3: 52–64.
- Yao T, Qin D, Shen Y, Zhao L, Wang N, Lu A. 2013. Cryospheric changes and their impacts on regional water cycle and ecological conditions in the Qinghai–Tibetan Plateau. *Chin. J. Nat. Med.* 35: 179–186.
- Yin Y, Wu S, Chen G, Dai E. 2010. Attribution analyses of potential evapotranspiration changes in China since the 1960s. *Theor. Appl. Climatol.* **101**: 19–28.
- Yin Y, Wu S, Zhao D, Zheng D, Pan T. 2013. Modeled effects of climate change on actual evapotranspiration in different eco-geographical regions in the Tibetan Plateau. J. Geogr. Sci. 23: 195–207.
- You Q, Kang S, Flügel W-A, Sanchez-Lorenzo A, Yan Y, Huang J, Martin-Vide J. 2010a. From brightening to dimming in sunshine duration over the eastern and central Tibetan Plateau (1961–2005). *Theor. Appl. Climatol.* **101**: 445–457.
- You Q, Kang S, Pepin N, Flügel W-A, Sanchez-Lorenzo A, Yan Y, Zhang Y. 2010b. Climate warming and associated changes in atmospheric circulation in the eastern and central Tibetan Plateau from a homogenized dataset. *Global Planet. Change* 72: 11–24.
- You Q, Kang S, Flügel WA, Sanchez-Lorenzo A, Yan Y, Huang J, Martin-Vide J. 2010c. Frombrightening to dimming in sunshine duration over the eastern and central Tibetan Plateau (1961–2005). *Theor. Appl. Climatol.* **101**: 445–457.
- You Q, Fraedrich K, Min J, Kang S, Zhu X, Pepin N, Zhang L. 2014. Observed surface wind speed in the Tibetan Plateau since 1980 and its physical causes. *Int. J. Climatol.* 34: 1873–1882.
- You Q, Min J, Jiao Y, Sillanpää M, Kang S. 2016. Observed trend of diurnal temperature range in the Tibetan Plateau in recent decades. *Int. J. Climatol.* 36: 2633–2643.
- Yu H, Luedeling E, Xu J. 2010. Winter and spring warming result in delayed spring phenology on the Tibetan Plateau. *Proc. Natl. Acad. Sci. U.S.A.* 107: 22151–22156.

- Yu J, Zhang G, Yao T, Xie H, Zhang H, Ke C, Yao R. 2016. Developing daily cloud-free snow composite products from MODIS Terra–aqua and IMS for the Tibetan Plateau. *IEEE Trans. Geosci. Remote Sens.* 54: 2171–2180.
- Zhang T. 2007. Perspectives on environmental study of response to climatic and land cover/land use change over the Qinghai–Tibetan Plateau: an introduction. *Arct. Antarct. Alp. Res.* **39**: 631–634.
- Zhang Y, Li T, Wang B. 2004. Decadal change of the spring snow depth over the Tibetan Plateau: the associated circulation and influence on the East Asian summer monsoon. *J. Climatol.* **17**: 2780–2793.
- Zhang X, Ren Y, Yin ZY, Lin Z, Zheng D. 2009. Spatial and temporal variation patterns of reference evapotranspiration across the Qinghai–Tibetan Plateau during 1971–2004. J. Geophys. Res. Atmos. 114: D15105.
- Zhang B, Wu Y, Zhu L, Wang J, Li J, Chen D. 2011a. Estimation and trend detection of water storage at Nam Co Lake, central Tibetan Plateau. J. Hydrol. 405: 161–170.
- Zhang G, Xie H, Duan S, Tian M, Yi D. 2011b. Water level variation of Lake Qinghai from satellite and in situ measurements under climate change. J. Appl. Remote Sens. 5(1): 053532.
- Zhang G, Xie H, Yao T, Liang T, Kang T. 2012. Snow cover dynamics of four lake basins over Tibetan Plateau using time series MODIS data (2001–2010). *Water Resour. Res.* **48**: W10529.
- Zhang D, Huang J, Guan X, Chen B, Zhang L. 2013a. Long-term trends of perceptible water and precipitation over the Tibetan Plateau derived from satellite and surface measurements. J. Quant. Spectrosc. Radiat. Transfer 122: 64–71.
- Zhang L, Su F, Yang D, Hao Z, Tong K. 2013b. Discharge regime and simulation for the upstream of major rivers over Tibetan Plateau. J. Geophys. Res. Atmos. 118: 8500–8518.
- Zhang Y, Cheng G, Li X, Han X, Wang L, Li H, Flerchinger GN. 2013c. Coupling of a simultaneous heat and water model with a distributed hydrological model and evaluation of the combined model in a cold region watershed. *Hydrol. Processes* 27: 3762–3776.
- Zhang C, Tang Q, Chen D. 2017a. Recent changes in the moisture source of precipitation over the Tibetan Plateau. *J. Clim.* **30**: 1807–1819. https://doi.org/10.1175/JCLI-D-15-0842.1.
- Zhang G, Yao T, Bolch T, Xie H, Chen D, O'Reilly CM, Yi S, Shum CK, He Y, Lei W, Shang K. 2017b. Climate change drives extensive lake area changes on Asia's high plateaus. *Geophys. Res. Lett.* 44: 252–260. https://doi.org/10.1002/2016GL072033.
- Zhang G, Yao T, Shum CK, Yi S, Yang K, Xie H, Feng W, Bolch T, Wang L, Behrangi A, Zhang H, Wang W, Xiang Y, Yu J. 2017c. Lake volume and groundwater storage variations in Tibetan Plateau's endorheic basin. *Geophys. Res. Lett.* 44: 5550–5560.
- Zhao L, Ping C, Yang D, Cheng G, Ding Y, Liu S. 2004. Changes of climate and seasonally frozen ground over the past 30 years in Qinghai–Xizang (Tibetan) Plateau, China. *Global Planet. Change* 43: 19–31.
- Zhao L, Wu Q, Marchenko S, Sharkhuu N. 2010. Thermal state of permafrost and active layer in central Asia during the international polar year. *Permafrost Periglacial Processes* **21**: 198–207.
- Zhu L, Xie M, Wu Y. 2010. Quantitative analysis of lake area variations and the influence factors from 1971 to 2004 in the Nam Co basin of the Tibetan Plateau. *Chin. Sci. Bull.* 55: 1294–1303.